



HEALTHCARE SAFETY
INVESTIGATION BRANCH

WWW.HSIB.ORG.UK



Clinical decision making: diagnosis and treatment of pulmonary embolism in emergency departments

Independent report by the
Healthcare Safety Investigation Branch I2019/016

March 2022

Providing feedback and comment on HSIB reports

At the Healthcare Safety Investigation Branch (HSIB) we welcome feedback on our investigation reports. The best way to share your views and comments is to email us at enquiries@hsib.org.uk or complete our online feedback form at www.hsib.org.uk/tell-us-what-you-think.

We aim to provide a response to all correspondence within five working days.

This document, or parts of it, can be copied without specific permission providing that the source is duly acknowledged, the material is reproduced accurately, and it is not used in a derogatory manner or in a misleading context.



About HSIB

We conduct independent investigations of patient safety concerns in NHS-funded care across England. Most harm in healthcare results from problems within the systems and processes that determine how care is delivered. Our investigations identify the contributory factors that have led to harm or the potential for harm to patients. The safety recommendations we make aim to improve healthcare systems and processes, to reduce risk and improve safety.

We work closely with patients, families and healthcare staff affected by patient safety incidents, and we never attribute blame or liability.

Considerations in light of coronavirus (COVID-19)

A number of national reports were in progress when the COVID-19 pandemic significantly affected the UK in 2020 and 2021. Much of the work associated with developing the reports necessarily ceased as HSIB's response was redirected.

For this national report, the investigation was initially paused, but then restarted due to its association with COVID-19. The processes HSIB used to engage with staff and families had to be adapted. Changes are described further in this report.

A note of acknowledgement

HSIB would like to thank the many healthcare staff who contributed their time and insights to the investigation, particularly during the added pressures of the COVID-19 pandemic. Their input has helped shape this report and allowed an in-depth exploration of decision making in clinical practice.

This report also shares the story of the care of a patient, Martin. HSIB did not specifically investigate the events affecting Martin, but his case helps convey the human and emotional importance of supporting improvements in patient safety. HSIB is grateful to Martin's Wife for sharing his experience.

About this report

This report is intended for healthcare organisations, healthcare staff, policymakers and the public to help improve patient safety in relation to the decisions made in the diagnosis and treatment of pulmonary embolism in emergency departments. The report focuses on clinical decision making and the challenges staff face when making decisions with limited information and time. For readers less familiar with the areas explored in this report, terms are explained throughout.

Since agreement of the safety recommendations outlined in this report plans were announced for Health Education England to merge with NHS England and NHS Improvement. While this does not change the recommendation made, the recipient of the recommendation may require updating in the future.

Our investigations

Our investigators and analysts have diverse experience of healthcare and other safety-critical industries and are trained in human factors and safety science. We consult widely in England and internationally to ensure that our work is informed by appropriate clinical and other relevant expertise.

We undertake patient safety investigations through two programmes:

National investigations

Concerns about patient safety in any area of NHS-funded healthcare in England can be referred to us by any person, group or organisation. We review these concerns against our investigation criteria to decide whether to conduct a national investigation. National investigation reports are published on our website and include safety recommendations for specific organisations. These organisations are requested to respond to our safety recommendations within 90 days, and we publish their responses on our **website**.

Maternity investigations

We investigate incidents in NHS maternity services that meet criteria set out within one of the following national maternity healthcare programmes:

- Royal College of Obstetricians and Gynaecologists' 'Each Baby Counts' report
- MBRRACE-UK 'Saving Lives, Improving Mothers' Care' report.

Incidents are referred to us by the NHS trust where the incident took place, and, where an incident meets the criteria, our investigation replaces the trust's own local investigation. Our investigation report is shared with the family and trust, and the trust is responsible for carrying out any safety recommendations made in the report.

In addition, we identify and examine recurring themes that arise from trust-level investigations in order to make safety recommendations to local and national organisations for system-level improvements in maternity services.

For full information on our national and maternity investigations please **visit our website**.

Executive Summary

This investigation was launched during the early stages of the COVID-19 pandemic, which meant that HSIB had to adapt its usual investigation method. Rather than investigating a specific patient safety incident, this investigation took an exploratory approach. It reviewed incidents associated with the diagnosis and treatment of pulmonary embolism in emergency departments to identify the factors that contributed to these incidents. The factors were then grouped into themes to identify the broader focus for a national investigation. One of the themes that emerged was clinical decision making, and this became the focus of the national investigation.

It is recognised that further factors will also influence the diagnosis and treatment of patients, and that patients with pulmonary embolism will seek help from other areas of healthcare. However, these issues were outside of the scope of this investigation.

Background

A pulmonary embolism (PE) is a blockage in the arteries of a person's lung, usually caused by a blood clot. Patients with a PE may have a range of symptoms (health problems that the patient experiences) and signs (things that clinicians find through examinations and tests). These include the classic symptoms of PE such as breathlessness and chest pain. PE may be difficult to diagnose and, if not recognised, can cause significant harm.

The purpose of this investigation is to identify learning that could support clinicians to make decisions about the diagnosis and treatment of PE patients in emergency departments (EDs). This learning may also apply to the treatment of other conditions where patients have non-specific symptoms.

Clinicians diagnose a PE by assessing a person's symptoms and signs, and then requesting appropriate tests. The decisions made during the diagnosis and treatment of PE can be influenced by several factors including the expertise of the staff who assess the patient, and the context within which they are working.

The investigation

HSIB identified a safety risk related to delayed or missed diagnosis and treatment of PE across several areas of healthcare where patients may first seek help. A review of events reported into the Strategic Executive Information System (a national database of patient safety incidents) identified EDs as one of these areas.

At the time of investigation access to EDs was limited because of the COVID-19 pandemic, and the investigatory approach was therefore adapted. Rather than examining a specific event, the investigation reviewed 14 serious incident reports obtained from hospitals across England. This review identified the terms of reference for a national investigation which aimed to:

- examine clinical decision making in the diagnosis and treatment of PE and the role of expertise (significant knowledge and skill that supports effective and practical decision making) using an Applied Cognitive Task Analysis (ACTA)
- identify factors in the wider healthcare system that support or inhibit diagnostic decision making when staff are seeing patients with non-specific symptoms and signs that may suggest PE.

This investigation was a first opportunity for HSIB to apply a specific methodology (ACTA) to examine decision making in healthcare.

Findings

- Recognising that a person may have a PE is challenging, particularly for less experienced staff and when the person's signs and symptoms are non-specific or atypical.
- Deciding whether to initiate treatment for a suspected PE requires a decision that balances risks, and this decision can benefit from expert knowledge and skill.
- Despite expertise and the available tools to help identify patients who may have a PE, a small number of PEs may always be missed.
- Experts use different thought processes and show different behaviours when making decisions compared to more novice staff.
- Decision-making skills in healthcare are commonly developed through experience, without formal training or opportunities to practise making decisions.
- Simulation-based learning has the potential to help staff acquire decision-making skills more quickly.
- Other industry sectors, such as aviation and the fire service, aim to accelerate the development of decision-making skills through structured training and the use of 'generic decision tools' for analytical decisions.
- EDs do not always provide the conditions which support the development of decision-making skills.

- Decision making in EDs is affected by workload, workforce availability, and performance targets.
- ED staff asked for further guidance to be provided on the use of decision aids to support the diagnosis of PE.
- The design of ED processes influences the decisions staff make. There is no standard model of initial patient assessment in EDs; this contributes to variation in the requesting of tests which can affect later decisions.
- Pathways for the diagnosis and treatment of PE in outpatient settings may create a safety risk where patients are discharged on anticoagulation medicines without a confirmed diagnosis; the capacity of imaging services is a significant contributor to this.
- Loss of clinical information when a patient's care is handed over was identified as a further safety risk. This can contribute to harm if tests, such as D-dimer (a blood test used as part of the assessment of likelihood of PE), are not followed up.
- Work procedures for the diagnosis and treatment of PE are not routinely designed in line with human factors principles to support their access and use.
- The physical design of environments may also affect decision making.

Safety recommendations and safety observations

HSIB wishes to acknowledge the challenges staff face in EDs when caring for patients with complex health conditions in the context of limited resource and capacity. This has been further exacerbated by the COVID-19 pandemic. HSIB recognises that these challenges mean the safety recommendations and safety observations in this report will be difficult to implement at the time of publication. Publication has gone ahead to ensure learning is shared and for the findings to be acted upon when capacity allows.

HSIB makes the following safety recommendations

Safety recommendation R/2022/188:

HSIB recommends that Health Education England works with appropriate professional bodies to develop and implement a strategy for supporting the education and training of clinical practitioners that can facilitate the development of decision-making skills. This strategy should consider the use of innovative approaches such as simulation and immersive learning.

Safety recommendation R/2022/189:

HSIB recommends that the National Institute for Health and Care Excellence reviews the findings of this investigation in relation to its guidance NG158, 'Venous thromboembolic diseases: diagnosis, management and thrombophilia testing', and updates the guidance if required.

Safety recommendation R/2022/190:

HSIB recommends that the Royal College of Emergency Medicine promotes best practice around diagnostic decision making with respect to patients with potential symptoms and signs of pulmonary embolism.

HSIB makes the following safety observations**Safety observation O/2022/155:**

It may be beneficial for healthcare to learn from other industries and develop its own evidence base on strategies to accelerate the development of expert decision-making skills. These strategies may include:

- development of a generic decision tool for implementation in healthcare training and clinical practice to support analytical decision making
- incorporation into education programmes of theory around how people make decisions and influences on decision making
- the use of simulation as a regular intervention to support practice and development of decision-making skills across scenarios with different levels of complexity
- consideration of the role of simulation in competency assessments for key skills.

Safety observation O/2022/156:

It may be beneficial if the findings of this investigation are used to support the development of staff expertise in decision making through:

- building understanding of how experts think and make decisions
- supporting reflection on the outcomes of simple and complex decisions
- development of clinical supervision skills of senior staff
- regular multidisciplinary case review.

Safety observation O/2022/157:

It may be beneficial for individual organisations to understand the extent to which national guidance on the diagnosis and management of pulmonary embolism is implemented across their organisations. This would help to identify local barriers to implementation to address. In particular it may be helpful to consider, in line with the findings of this investigation, local engagement with the scoring systems available to help predict the likelihood of a pulmonary embolism.

Safety observation O/2022/158:

It may be beneficial for emergency departments and same-day emergency care units to have rapid access to recommended imaging for patients who require it for the diagnosis of pulmonary embolism.

Safety observation O/2022/159:

It may be beneficial for the positivity standard for computerised tomography pulmonary angiography (CTPA) (that at least 15% of CTPAs should show a pulmonary embolism) to be evaluated to understand its effects on emergency department decision making.

Safety observation O/2022/160:

It may be beneficial for healthcare work procedures to be written in line with the principles for effectiveness and usability provided by the Chartered Institute of Ergonomics and Human Factors.

Contents

- 1 Background and context **12**
- 2 Involvement of the Healthcare Safety Investigation Branch **23**
- 3 Local factors – review of serious incidents **29**
- 4 National investigation analysis and findings **35**
- 5 Summary of findings, safety recommendations and safety observations **68**
- 6 References **72**
- 7 Appendices **82**

1 Background and context

1.1 Pulmonary embolism

1.1.1 Venous thromboembolism (VTE) is a collective term for the formation of blood clots (thrombus) which can travel through and then block (occlude) a person's blood vessels. This process is known as embolism. VTE includes deep vein thrombosis (DVT, blood clots usually in the lower leg/calf) and pulmonary embolism (PE). A PE is a blockage in the arteries of a person's lung usually caused by a blood clot.

1.1.2 The incidence (occurrence rate) of PE per year is around 60 to 70 cases per 100,000 people (Howard et al, 2018). The individual risk of a person developing a VTE is influenced by their risk of forming blood clots.

PE risk factors and signs/symptoms

1.1.3 A PE is the partial or complete blockage of single or multiple pulmonary (lung) blood vessels, usually from a DVT that travels from a person's leg to their lung. There are multiple factors that increase the risk of a person developing a PE, these include (Clarity Informatics, 2020):

- cancer
- immobility
- inflammation including infection
- inherited disorders of blood clotting
- major injury or injury to the lower limbs
- obesity
- oestrogen containing medicines
- pregnancy
- previous VTE in the individual or their family
- surgery, particularly bone and emergency.

1.1.4 Clinicians use symptoms (health problems that the patient experiences) and signs (things that clinicians find through examinations and tests) to help them make a diagnosis. The classic symptoms of PE (see table 1) include pleuritic chest pain (pain that is worse on breathing in), shortness of breath and haemoptysis (coughing up blood). However, patients may also show symptoms and signs that are few, unexpected (atypical), or do not point at a particular diagnosis (non-specific). Patients may also arrive at an emergency department following a collapse and cardiac arrest. This variation can make diagnosis challenging.

Table 1 Examples of symptoms and signs of a pulmonary embolism adapted from the PE clinical knowledge summary (Clarity Informatics, 2020)

Symptoms suggestive of PE:	Shortness of breath, fast heart rate, chest pain worse on breathing in and/or features of DVT
Other symptoms may include:	Chest pain behind the breastbone, cough, coughing up blood, dizziness and collapse
Other signs may include:	Low oxygen levels, abnormal heart rhythm and low blood pressure

Diagnosing PE

- 1.1.5 The National Institute for Health and Care Excellence (NICE) provides guidance on the diagnosis and treatment of adults with PE (National Institute for Health and Care Excellence, 2020a). NICE states that patients seeing a clinician with symptoms and/or signs of PE should have a full medical history, physical examination and chest X-ray.
- 1.1.6 Where there is a suspicion of PE, NICE recommends the use of a two-level Wells score (**see table 2**). The Wells score is a validated tool to help predict the likelihood of a PE when used in conjunction with a D-dimer blood test (see 1.1.7). Where the Wells score is more than four points, PE is likely and patients should be offered radiological imaging (a scan). Where the Wells score is four or less, PE is unlikely and a D-dimer blood test can be considered.
- 1.1.7 D-dimer is a protein that is made when a blood clot breaks down. A D-dimer blood test checks for the levels of this protein in the blood. This test is widely available in hospitals and results are usually available within a few hours. A negative D-dimer test result suggests no PE is present in a patient who is thought to be unlikely to have a PE (based on Wells score). If the D-dimer is positive, NICE recommends that the patient has a scan to look for a PE (National Institute for Health and Care Excellence, 2020a).

Table 2 Two-level PE Wells score (National Institute for Health and Care Excellence, 2020a)


Clinical feature	Points
Clinical signs and symptoms of DVT (minimum of leg swelling and pain with palpation [examination by touch] of the deep veins)	3
An alternative diagnosis is less likely than PE	3
Heart rate more than 100 beats per minute	1.5
Immobilisation for more than 3 days or surgery in the previous 4 weeks	1.5
Previous DVT/PE	1.5
Haemoptysis [coughing up blood]	1
Malignancy [cancer] (on treatment, treated in the last 6 months, or palliative [non curable])	1

1.1.8 D-dimers are not specific to PE. A person's D-dimer levels can also be raised by factors such as age, ethnicity, pregnancy, recent surgery, certain bleeding disorders, cancer, heart disease, trauma, aortic blood vessel problems and COVID-19.

1.1.9 The main imaging technique used to support the diagnosis of PE is computerised tomography pulmonary angiography (CTPA). CTPA provides images of a patient's pulmonary (lung) blood vessels. Where filling of blood vessels is not seen, this can suggest a PE (**see figure 1**). The ventilation (V)/perfusion (Q) scan is an alternative for certain patients who cannot or should not have a CTPA (National Institute for Health and Care Excellence, 2020a). V/Q scans were not considered in this investigation.

Figure 1 CTPA images showing a possible pulmonary embolism (Choong et al, 2008)

Darker area in the 'white' of the blood vessel shows reduced filling of the blood vessel suggesting a PE



Treating PE

1.1.10 Anticoagulation medicines are the main treatment for suspected and confirmed PE. These medicines reduce the formation of further blood clots. Due to the risk of clots increasing in size while patients are undergoing tests for suspected PE, NICE recommends interim treatment with anticoagulation in certain circumstances (National Institute for Health and Care Excellence, 2020a):

- if a patient has a likely PE and cannot have an immediate scan
- if a patient is unlikely to have a PE (they have a Wells score of four or less) and a D-dimer result cannot be obtained within 4 hours.

1.1.11 After a diagnosis of PE is confirmed, patients continue treatment with longer-term anticoagulant medicines. Other treatments for life-threatening PEs include medicines to break down clots (thrombolytics) and procedures to directly remove clots.

Outpatient diagnosis and treatment of PE

- 1.1.12 Traditionally patients with a suspected PE would have been admitted to hospital. More recently there has been a move to managing many conditions outside of hospital when patients are well enough to be at home. This improves patient experience and reduces demand on hospitals.
- 1.1.13 In 2018 the British Thoracic Society (BTS) developed guidance for the initial treatment of patients with PE as outpatients (Howard et al, 2018). NICE recommends considering 'outpatient treatment for suspected or confirmed low-risk PE' and recommends using a validated tool to determine patients who are low risk of 'poor outcomes' from a PE (National Institute for Health and Care Excellence, 2020a). The BTS provides suggestions on what validated tools can be used (Howard et al, 2018).

COVID-19 and PE

- 1.1.14 During the COVID-19 pandemic it became apparent that the virus affected the formation of blood clots. COVID-19 can lead to patients developing PE, and other clots, through immune-related clot formation. The virus can also result in increased D-dimers in the blood because of inflammation.
- 1.1.15 Studies have explored the presence of PE in COVID-19 with variable results. These have been summarised in various reports and systematic reviews (for example, Tholin et al, 2021; Jiménez et al, 2021; Malas et al, 2020). Clot formation is likely related to the severity of a patient's COVID-19 infection.
- 1.1.16 The symptoms that a patient may have with COVID-19 may be similar to those of a PE. COVID-19 patients may be breathless and have a cough. This can make distinguishing between the diagnoses difficult. As described above, patients may also have COVID-19 and a PE at the same time.

1.2 Emergency departments

- 1.2.1 This investigation is focused on type-one emergency departments (EDs). Type-one EDs are consultant-led, available 24 hours a day and can treat any emergency patient. As of 2019/20 there were 134 hospital trusts with type-one EDs in England (NHS Digital, n.d.).
- 1.2.2 In England, at the time of writing, most EDs are directed by a national operating standard under which 95% of people attending an ED should be admitted to hospital, transferred or discharged within 4 hours of arrival (NHS England and NHS Improvement, 2020). In December 2021 there were 1,250,669 attendances at major EDs in England with 73.3% of attendances meeting the national operating standard (NHS England and NHS Improvement, n.d.).



ED processes


- 1.2.3 When patients arrive at an ED, staff will first prioritise and direct them to the most appropriate place for treatment. This happens through an 'initial assessment' process (**see figure 2**) and triage. Initial assessment aims to rapidly identify those patients who require immediate treatment, and also helps to ensure patients are seen and treated quickly.
- 1.2.4 Following initial assessment, patients will be diagnosed and treated, and then discharged or referred on for specialist care. Patients may be seen by an ED doctor, advanced clinical practitioner (ACP) or general practitioner (GP) in the ED.

PE pathways

- 1.2.5 To support the standardised assessment of patients with different symptoms, EDs have clinical pathways and work procedures. These include pathways and procedures for the diagnosis and treatment of PE, usually based on NICE guidance.
- 1.2.6 PE pathways include a combination of tools to support diagnostic decision making (such as Wells score), D-dimer blood testing and radiological imaging (scans). PE pathways support the standardised evaluation of patients and reduce the risk of subsequent death from PE (van Belle et al, 2006). These pathways will not identify every patient with a PE, but will help identify the majority.



Figure 2 Example of the initial assessment process for the diagnosis and treatment of ED patients, adapted from the Royal College of Emergency Medicine (2017).



1.3 Decision making

- 1.3.1 Decision making is ‘the process of reaching a judgment, choosing an option or course of action to meet the needs of a given situation’ (Clinical Human Factors Group, 2018). In this report, the phrase ‘clinical decision making’ is used to describe how healthcare staff seek information, judge information, decide on possible diagnoses and treatment, and implement a plan.
- 1.3.2 Decision making has been extensively researched and traditionally is seen to involve two approaches (Kahneman, 2012):
- System 1 (fast): recognising patterns and making rapid and intuitive decisions.
 - System 2 (slow): developing ideas for what may be needed for less recognisable situations and testing it by gathering and analysing information (also referred to as analytical decision making).
- 1.3.3 A person’s decisions are influenced by their recognition of and familiarity with a situation. Unfamiliar situations are more likely to result in slower, analytical thinking to reach a decision (system 2). When a situation is recognised, rapid and intuitive decisions (system 1) are more likely to follow, supported by unconscious mental shortcuts, known as ‘heuristics’. Heuristics support decision making in situations where there is limited time or information (an example is provided in figure 3).

Figure 3 Example of a heuristic (Clinical Human Factors Group, 2018)

‘Selecting salbutamol nebuliser and oxygen as first line treatment for wheeze, as it is the most common treatment administered.’
(Clinical Human Factors Group, 2018)

- 1.3.4 Heuristics are useful because they can help staff reach a diagnosis and start treatment quickly. However, if the outcome of the heuristic is not as intended, with hindsight it may be called a bias (Klein, 1989). Bias is often the result of limited information and limited prompts.
- 1.3.5 More recently, the traditional beliefs around clinical decision making have been challenged because they may not account for how staff make decisions in the ‘real world’. Newer models incorporate the role of intuition, the influence of emotions, self-regulation and the real-world context (Patel et al, 2015).

Recognition-primed decision making

- 1.3.6 An individual's clinical decision making is influenced by their knowledge, experience and memory, and the situation within which they find themselves. Naturalistic decision making (NDM) refers to how people make decisions in complex and real-world situations (Klein, 2008). Research into NDM led to the development of the 'recognition-primed decision making' (RPD) model.
- 1.3.7 The RPD model was developed to help understand how experts make decisions in complicated, information limited, and/or time-critical situations (Klein, 1997). An expert is someone who has significant knowledge about and skill in a particular subject or area, leading to high-quality performance.
- 1.3.8 When confronted with a situation, experts will often recognise the situation leading to a course of action. RPD helps describe how experts recognise, make sense of, and identify an appropriate course of action through three variations depending on the situation (Helander, 2006; Klein, 1999):
- Simple match – the person experiences a situation that they perceive as familiar and typical, resulting in a familiar course of action.
 - Sensemaking required – the person experiences a situation which they interpret to be slightly different from what they are familiar with or from what they expected, resulting in gathering further information and engaging in further assessment before implementing a familiar course of action.
 - Evaluate the action – the person experiences a familiar situation, but there are factors that mean they cannot implement the familiar course of action. The course of action is modified and the person mentally evaluates or simulates whether it will work before implementing it.

Table 3 provides some real-world examples.



Table 3 Examples of the variations of recognition-primed decision making


Variation	Example
Simple match	The consultant sees a patient with pleuritic chest pain, shortness of breath and a swollen leg. They recognise that the patient likely has a PE and implement appropriate treatment.
Sensemaking required	The consultant sees a patient with pleuritic chest pain and shortness of breath. They ask questions to get more information, including risk factors for blood clotting. They identify that the patient likely has a PE and implement appropriate treatment.
Evaluate the action	The consultant sees a patient with pleuritic chest pain, shortness of breath and a swollen leg. They recognise that the patient likely has a PE and plans to start treatment with anticoagulation in line with their hospital's policy. The patient is allergic to the anticoagulation that the policy advises and so an alternative medicine is used following consideration of the risks and benefits.

Expert versus novice decision makers

- 1.3.9 Expert decision makers are those with significant knowledge and skill that support effective and practical decision making. Expertise is different to experience, which relates to the acquiring of knowledge and skill over years. Years of experience do not necessarily mean someone is an expert, but it can contribute to the development of expertise.
- 1.3.10 To develop decision making as a skill, practice is required. This practice includes developing appropriate heuristics for rapid decision making, and using structured approaches to analytical decision making. Skill development progresses through levels from novice to expert (**see figure 4**). More novice or beginner decision makers may be knowledgeable, but have limited real-world practice; they therefore need to be supervised.
- 1.3.11 Novice decision makers tend to default to following rules, meaning a situation is matched with a rule to help decide on an action. Checklists and procedures (cognitive aids) support rule-based decisions. While rule-based decision making allows rapid action, there is a risk that rules can be misapplied, or that no rule can be found to match a situation. In complex healthcare situations not all situations can be anticipated and therefore analytical decision making is required.

1.3.12 More expert decision makers have a broader mental repertoire to help them deal with situations. They are therefore able to deal with more complexity. Because of increased practice and experience, their heuristics are less prone to being misapplied. They may also be more able to identify when a situation is outside of their mental repertoire and requires practiced analytical decision making.

Figure 4 Model of skill acquisition adapted from Dreyfus (2004)



2 Involvement of the Healthcare Safety Investigation Branch

This section outlines how HSIB became aware of the safety risks surrounding the timely diagnosis and treatment of pulmonary embolism (PE) in emergency departments (EDs). It also describes the actions undertaken by the investigation, including the review of incidents in England to identify areas on which to focus a national investigation.

2.1 Decision to investigate

- 2.1.1 Following HSIB's investigation into the 'Management of venous thromboembolism risk in patients following thrombolysis for an acute stroke' (**Healthcare Safety Investigation Branch, 2020a**), a further safety risk was identified of delay in the recognition of PE.
- 2.1.2 HSIB reviewed the Strategic Executive Information System (StEIS, the national reporting system for serious incidents) for other incidents involving a delay in the diagnosis of PE (**see appendix 7.1** for the search criteria used). This identified several incidents, particularly in EDs, before and during the COVID-19 pandemic.
- 2.1.3 As a result of this information, it was recommended to HSIB's Chief Investigator that a national investigation should be undertaken. The safety risk met the following criteria for a national investigation.

Outcome impact - what was, or is, the impact of the safety issue on people and services across the healthcare system?

- The incidence (occurrence rate) of PE per year is around 60 to 70 cases per 100,000 people (Howard et al, 2018). While overall trends may be decreasing, it presents a significant burden on public health services (Barco et al, 2020).
- PE is a significant condition associated with a risk of death (Heit et al, 2016). PE can also result in other complications including heart rhythm abnormalities (arrhythmias), heart failure, and lung tissue damage (pulmonary infarction).
- If undiagnosed or untreated, PE is associated with higher risks of death or long-term harm. Incidents reported on StEIS demonstrate the continuing harm associated with delays to PE diagnosis and its treatment.

- PE has a significant psychological impact on patients, families and staff. Patients with a PE may have psychological distress (for example, Tran et al, 2021). Healthcare staff who may have missed a PE when diagnosing a patient may also experience distress (**Healthcare Safety Investigation Branch, 2021a**).

Systemic risk – how widespread and how common a safety issue is this across the healthcare system?

- In 2019 the National Confidential Enquiry into Patient Outcome and Death (NCEPOD) identified delays in the processes for treating patients with PE in almost 40% of the 526 cases it reviewed (National Confidential Enquiry into Patient Outcome and Death, 2019).
- NCEPOD also identified that for just under 20% of patients there was a delay in starting anticoagulation medication, and there was variation in how patients were selected for outpatient care pathways.
- The incidents reported on StEIS demonstrated a risk in EDs. ED environments may be higher risk because patients may visit the ED with non-specific symptoms.

Learning potential – what is the potential for an HSIB investigation to lead to positive changes and improvements to patient safety across the healthcare system?

- The data demonstrates a need to understand why delays to diagnosis and treatment of PE occur in EDs and how they can be mitigated. An investigation offers the potential to understand contributory factors.
- COVID-19 may also impact on the diagnosis of PE (**see 1.1.14**). Understanding the contributory factors in delays to diagnosis and treatment of PE may add value to the treatment of PE in patients with COVID-19.

2.2 Evidence gathering and investigation approach

Investigative approach

2.2.1 The investigation was carried out between October 2020 and May 2021. The investigation had two parts:

- 1 Review of serious incidents (local factors): a review of serious incidents in England to identify local factors that contributed to incidents related to diagnosis and treatment of PE in EDs. The findings defined the terms of reference for part two.

- 2 National investigation: a national investigation into the factors that influenced ED staff decision making in the diagnosis and treatment of PE.

Review of serious incidents (local factors)

2.2.2 The investigation approached NHS hospitals to request copies of serious incident investigation reports where the incident:

- occurred during 2020 in the NHS in England
- involved ED and the diagnosis and/or treatment of PE, and
- was associated with delays and/or missed diagnosis and/or treatment.

2.2.3 The investigation analysed the serious incident reports to identify the factors that contributed to the incidents. The reports were analysed using the Systems Engineering Initiative for Patient Safety (SEIPS) (Holden et al, 2013; Carayon et al, 2006) as a framework (**see appendix 7.2**). The approach was similar to that of other HSIB reports (**Healthcare Safety Investigation Branch, 2021b**). Findings are presented in **section 3**.

Terms of reference for national investigation

2.2.4 The serious incident reports demonstrated various factors that influenced the occurrence of the incidents. A consistent finding across the serious incident investigations was that 'incorrect' decisions were made in the diagnosis and treatment of PE. The aims of the national investigation were therefore to attempt to understand the nuances of the decisions made in EDs and factors that could affect them. The terms of reference were to:

- examine clinical decision making in the diagnosis and treatment of PE and the role of expertise
- identify factors in the wider healthcare system that support or inhibit diagnostic decision making when staff are seeing patients with non-specific symptoms and signs that may suggest PE.

National investigation

Examining clinical decision making

2.2.5 Applied Cognitive Task Analysis (ACTA) was used to examine clinical decision making. ACTA is a technique to analyse the cognitive elements of a task. It considers how expertise supports the undertaking of cognitive tasks and why such tasks may be difficult for novices.

2.2.6 ACTA methodology is provided in a toolkit (Militello et al, 1997; Militello and Hutton, 1998) with guidance for use. The steps of an ACTA, and how the investigation undertook the analysis, are provided in appendix 7.3. The methodology includes three methods and the investigation used two of these in the cognitive task diagram (a visual representation of the tasks involving thinking) and knowledge audit (interviews with staff to explore expertise using probing questions). Knowledge audit interviews were undertaken with clinical experts and novices in EDs (**see table 4**).

2.2.7 The terms 'expert' and 'novice' were used in line with the original ACTA model. To distinguish between levels of expertise the investigation was guided by Dreyfus' model (Dreyfus, 2004) (**see section 1.3** and **figure 4**). Staff were considered to be either experts or novices:

- Experts – senior nursing staff; advanced clinical practitioners; senior doctors in emergency medicine, including consultants.
- Novices – students of any profession; newly qualified nurses, doctors and advanced clinical practitioners; and those with limited situational knowledge and skill in emergency medicine.

Identifying wider system factors

2.2.8 Alongside the ACTA, the investigation undertook observations and interviews (**see table 4**) to identify factors in the wider healthcare system that influenced decision making. Interviews were also undertaken with national bodies and subject experts. Evidence was further gathered from academic and policy literature.

2.2.9 The investigation visited six EDs across four hospital trusts in England. The EDs were geographically spread and of different sizes with between 150 and 650 attendances a day (**see table 4**).

Table 4 Sources of information for the national investigation

Area of interest	Examining decision making	Wider healthcare system
Emergency departments	Trust 1: ED 1 and ED 2 (400 to 550 attendances/day) Trust 2: ED 3 (550 attendances/day) Trust 3: ED 4 and ED 5 (150 to 450 attendances/day) Trust 4: ED 6 (650 attendance/day)	
Local stakeholders	Doctors in emergency and acute medicine (foundation trainees to consultants) Advanced clinical practitioners and physician associates Nursing staff (bands five to eight) Paramedics Students (nursing, medicine and physician associates)	Radiology and radiography staff Emergency and acute medicine department managers Patient safety leads
National stakeholders	Health Education England 825 Naval Air Squadron, Royal Navy Royal Navy Safety Centre Chief Fire Officer	NHS England and NHS Improvement Same Day Emergency Care Programme British Thoracic Society National Institute for Health and Care Excellence National expert in haemostasis and thrombosis Royal College of Radiology ‘Getting it Right First Time’ Lead for Emergency Medicine Royal College of Emergency Medicine General Medical Council Medical Schools Council
Subject matter advisors	Applied cognitive psychology (ACTA and decision making)	Consultant in Emergency Medicine Consultant Haematologist

Analysis

- 2.2.10 Interviews were undertaken using a semi-structured approach and observations using an observation template based on SEIPS. Field notes were taken and collated for analysis.
- 2.2.11 The learning, safety observations and safety recommendations in this report were shared with stakeholders. This allowed further corroboration of findings, adjustment where necessary, and formation of safety recommendations that have been agreed by relevant national bodies.

Limitations

- 2.2.12 The investigation's findings are specific to decision making in EDs. This was the focus of the investigation and where the lines of inquiry led. Evidence collection was predominantly qualitative through the interviews, observations and ACTA. The findings are therefore based on staff experiences, perceptions and narrative. However, they have been verified with national experts, subject matter advisors and relevant national bodies.
- 2.2.13 The investigation acknowledges that there will be other factors that influence decision making outside of those considered in this report and that similar challenges will be seen in other parts of healthcare. The findings of this report may have wider benefit for decision making outside of the ED.
- 2.2.14 Safety recommendations have been made where the investigation has appropriate evidence and following engagement with the relevant national bodies. Safety observations represent learning from the investigation that may be of benefit to organisations and the wider healthcare system.

3 Local factors – review of serious incidents

This section provides a summary of findings from the investigation’s review of serious incident reports obtained from hospitals. The reports were analysed to identify local factors that contributed to delayed or missed diagnosis or treatment of pulmonary embolisms in emergency departments. The factors were grouped to identify common themes across the reports.

In one of the serious incident reports, the investigation learned about the care of Martin. The investigation approached Martin’s Wife who described the events; they are detailed here to provide insights into a real event and share the importance of learning from incidents. HSIB did not specifically investigate Martin’s care.

3.1 Martin’s story

- 3.1.1 Martin was a husband and father of two. He was 51 years old, and described by his Wife as “clever, but humble, and well-liked by everybody”. He was active in the local community and also enjoyed playing cricket and golf. He had been working from home during the COVID-19 pandemic as his office had been closed.
- 3.1.2 Martin was rarely unwell, but while working from home on a Friday, he started feeling short of breath. Martin and his Wife thought he could have COVID-19, so booked a test which was undertaken on the Sunday. Two days later the test came back as negative. Martin was still breathless so he rang NHS 111 and was advised to go to the local emergency department (ED).
- 3.1.3 Martin attended the ED on his own because of COVID-19 restrictions. He had some tests and was discharged home. Martin told his Wife that he had undergone a heart tracing, chest X-ray and blood tests, but the doctors could find nothing wrong so thought it might be a chest infection.
- 3.1.4 Over the following days Martin remained short of breath, but was still able to get on with daily life. Eight days after going to the ED, Martin drove into work for a meeting, but when he returned to his car became unwell. An ambulance was called which took him to the ED. Martin’s Wife received a call from the hospital asking her to pack a bag for him because he was going to need to stay in hospital.
- 3.1.5 About 30 minutes later Martin’s Wife received another call from the hospital to tell her that Martin had become seriously unwell. She was asked to get to the hospital as soon as possible and arrived about 45 minutes later. Martin



had suffered a cardiac arrest and was being moved to intensive care. Martin did not recover and died later the same day. A post-mortem report showed Martin had died from a pulmonary embolism (PE).

3.1.6 An investigation was undertaken by the hospital to find out whether there had been an opportunity to identify a PE during Martin's first visit to the ED. The hospital found that a PE had been considered and a D-dimer blood test to rule-out a PE had been planned, but was not requested. Martin's care was verbally handed over to another member of staff at shift change. Martin was later discharged as all his tests were normal, but staff did not know that a D-dimer test had not been undertaken.

3.1.7 The hospital's investigation highlighted issues relating to:


- who orders a D-dimer blood test and when
- handover of outstanding tests required
- high workload
- lack of relatives in ED to provide a further information
- overcrowding of the ED.

3.2 Serious incident overview

3.2.1 The investigation reviewed 14 serious incident reports, referred to here as 'reports'. Each report investigated the care of a patient who had an incorrect, missed or delayed diagnosis or treatment of PE in an ED. Several of the reports described patients who had died as a result of a PE. Some patients died from other conditions alongside a PE, or died from a condition incorrectly suspected to be a PE.

3.2.2 The reports commonly considered how the process of decision making by staff resulted in delayed or missed diagnoses of PE, and explored the associated contributory factors to a greater or lesser degree. The reports also considered the process of developing treatment plans for patients with suspected PE. Figure 5 provides an overview of the processes and outcomes seen in the reports.

Figure 5 Processes and outcomes found in the investigation's review of serious incident reports



3.3 Factors affecting diagnostic decision making

- 3.3.1 Diagnostic decision making refers to how ED staff use information to conclude whether a patient may have a PE. Decision aids and tools may be used to support decision making; these aids and tools may be available in procedures and guidelines.
- 3.3.2 The reports included missed diagnoses of PE, including where an alternative diagnosis was suspected, and delayed diagnoses of PE. Where a PE was not initially suspected, but was subsequently found, alternative initial diagnoses included chest infections (including COVID-19), muscular pain and anxiety. Reports also included examples of patients who were suspected of having a PE, but an alternative diagnosis was later found, such as aortic dissection (a tear that allows blood to flow between the layers of the wall of a person's aorta, which is the largest artery in the body).
- 3.3.3 Several contributory factors were found to have influenced the decisions made by staff in the diagnosis of PE. The recurrent themes were:
- COVID-19 was perceived to have increased the risk of missing a PE because the infection caused similar symptoms and could occur at the same time as a PE. There was limited national guidance to help clinicians identify patients with a PE and COVID-19.
 - Demand on EDs and the national operating standard were described as creating 'pressure' which limited time to 'pause' and consider the range of possible diagnoses.
 - Staffing shortages nationally and locally of senior doctors limited opportunities for in-person reviews of patients meaning they were assessed by more novice staff.
 - ED diagnostic and treatment pathways did not always prompt clinicians to consider PE when patients had symptoms that were unexplained or not typical of PE. Outpatient PE pathways did not always consider the risk of discharging a patient with an unknown diagnosis who was on anticoagulation medication, while they waited to have a scan. Examples included a case where the patient was later found to have an aortic dissection.
 - Conflict between ED and radiology departments was described, creating barriers to requesting imaging. Scans were a limited resource.
 - Decision making tools, cognitive aids and risk assessments to support diagnosis of PE were often not accessed or used.



- D-dimer blood tests influenced consideration of PE. Where a D-dimer had been requested during the initial assessment of a patient, PE became the lead potential diagnosis and where a D-dimer was not requested, PE was given less consideration. An elevated D-dimer was assumed to be related to PE and did not lead staff to consider other causes, such as aortic dissection.
- Review of results of tests or observations outside expected ranges did not always take place before a patient was given a diagnosis or discharged. Findings that were outside of expected ranges were not always accounted for. Examples included the D-dimer and fast heart rates.
- Patient symptoms were not always typical of PE or were non specific. Other symptoms included nausea and vomiting, abdominal pain and dizziness. Patients often had risk factors for blood clots, but some of these may have not been recognised, such as being less active during COVID-19.

3.3.4 Most reports contained a focus on how staff ‘biases’ had resulted in PE being considered less likely or other diagnoses being considered more likely. Prompts influencing bias and leading staff to consider alternative diagnoses to a PE included COVID-19, tender chest walls (muscular-related diagnosis), and mental health histories (anxiety-related diagnosis). The focus of investigations on individual bias limits learning by reducing consideration of the wider healthcare system. Instead, investigations should seek to identify the conditions that contribute to bias, and address them to support effective decision making. This includes supporting staff to develop more sophisticated and appropriate heuristics (**see 1.3.3**).

3.4 Factors affecting treatment

- 3.4.1 The reports included examples of delays in giving anticoagulation medication, or anticoagulation being given incorrectly, to patients with suspected or confirmed PE. Reports also described limited imaging capacity that meant immediate scans were uncommon where patients were not unwell.
- 3.4.2 Underdosing of anticoagulation was seen where prescribing was based on the estimated weight of the patient. This safety risk was identified nationally in 2010 (National Patient Safety Agency, 2010). Reports described insufficient emphasis on weighing patients, poor weighing equipment, limited prompts to identify which patients need weighing, and difficulties weighing unwell patients.
- 3.4.3 Three reports also identified challenges with the administration of thrombolysis (a medicine to break down blood clots) for patients with a PE in emergency situations. Staff found barriers to accessing medicines



because of unclear local processes for access to emergency medicines, limited signposting of medicine locations, and local training that did not include how to access the medicines.

3.5 Summary

3.5.1 The investigation's review of the reports provided an understanding of local and some national factors that contributed to the serious incidents. The investigation recognises that the number of reported incidents was small compared to the number of patients who arrive at EDs with a suspected PE. Some of the reports also represented patients who did not have the typical symptoms of PE, which made diagnosis more challenging.

3.5.2 From the review of the reports, the investigation concluded that:

- Decision making in EDs is challenging, particularly in the diagnosis of suspected PE where patients may have non-specific or atypical symptoms and signs.
- Local investigations often attributed incorrect decisions to 'biases' with limited exploration of how staff made decisions in practice.
- Staff did not always use the tools available to them to help make decisions, such as decision aids and guidelines.
- There are specific, wider ED factors that potentially contribute to incorrect decisions such as demand on services, workload, process and procedure design, conflict between departments, and access to support.

3.5.3 The findings suggested that there would be value in the investigation seeking to understand, in more depth, how ED staff make decisions, how staff can be supported to make effective decisions, and the wider factors that may influence those decisions.



4 National investigation analysis and findings

The national investigation aimed to understand the nuances of the decisions made in EDs and factors that could affect them. The findings are described in the following sections:

- **Section 4.1** – introduces the investigation’s observations of decision making in the emergency departments (EDs) visited.
- **Section 4.2** – examines the investigation’s first term of reference – to examine clinical decision making in the diagnosis and treatment of pulmonary embolism (PE) in EDs – with the aim of understanding in more depth how ED staff make decisions.
- **Section 4.3** – examines, based on the findings of the Applied Cognitive Task Analysis (ACTA), how decision-making expertise could be better developed in healthcare staff, with associated safety observations and safety recommendations.
- **Section 4.4** – explores the investigation’s second term of reference – to identify wider factors in the healthcare system that support or inhibit diagnostic decision making when staff are seeing patients in EDs with non-specific symptoms and signs that may suggest PE.

4.1 Decision making in emergency departments

- 4.1.1 HSIB considered decision making in EDs in a previous investigation, ‘Delayed recognition of acute aortic dissection’ (**Healthcare Safety Investigation Branch, 2020b**). Similar to that previous investigation, this investigation found staff using different approaches to decision making depending on the situation, their expertise and experience.
- 4.1.2 This investigation identified examples of staff using recognition-primed decision making (RPD, **see 1.3.6 to 1.3.8**). This was particularly noted when EDs were busy, when there were lots of patients waiting, and in urgent situations. Seeing patients with familiar signs and symptoms resulted in familiar actions.
- 4.1.3 When staff saw patients with unfamiliar signs and symptoms, particularly those that were less urgent, they described a more analytical approach. The investigation observed staff thinking through symptoms and signs, talking out loud, and discussing options with colleagues. While this approach was more comprehensive, it was time consuming and sometimes still did not result in the intended outcomes.

- 4.1.4 During observations the demand on EDs was seen to increase during the day. As demand grew, the available time to gather information about patients was reduced. The use of RPD in these situations supported rapid decisions, but some decisions were based on limited information. There was a risk that heuristics (**see 1.3.3**) may be misapplied as demonstrated in the serious incidents in **section 3**.
- 4.1.5 Staff also told the investigation how they managed uncertainty. Decisions were commonly based on limited information and without a full awareness of the situation. Staff aimed to identify common diagnoses and exclude “worst-case scenarios” (life-threatening conditions).
- 4.1.6 Staff described how managing uncertainty meant taking risks. For example, they knew that they could not know everything about a patient’s medical history, and were not able to test every patient for a PE. This demonstrated the need to make ‘trade-offs’, balancing factors such as accuracy, efficiency and thoroughness.
- 4.1.7 More novice staff and students were also heard matching patients to textbook descriptions of diagnoses, sometimes using educational resources. The ACTA subject matter advisor described these as rule-based decisions which are common for novices, but may risk rules being inappropriately applied to less familiar, more complex or nuanced situations. For example, novices generally associated a raised D-dimer with a potential PE, not considering other potential diagnoses.
- 4.1.8 Students and doctors in their early years of practice recited classical textbook features of PE to the investigation. These were chest pain, breathlessness and coughing up blood (haemoptysis). The investigation asked experts about the signs and symptoms of PEs and heard that these classic features were rarely all present; often a patient would have breathlessness alone, or a fast heart rate. Patients could also have atypical signs and symptoms such as described in the serious incident reports. The research suggests that 97% of patients with PE have at least one of the three classic features (BMJ Best Practice, 2021).

4.2 Examining clinical decision making


The investigation used ACTA (**see 2.2.5 to 2.2.7**) to examine the key decisions made by ED staff relating to the diagnosis and treatment of PE. During the development of a cognitive task diagram (**see figure 6**), ED staff identified three cognitive tasks as the most challenging. They told the investigation that

these tasks were influential in the diagnosis and treatment of suspected PE and involved decisions that needed the most expertise. The tasks were:

- initial assessment of the patient, including selecting investigatory tests
- further assessment and review of preliminary test results
- prescribing of anticoagulation medication while waiting for the scans or results of tests (referred to as 'interim coagulation'), particularly if the patient is to be discharged for outpatient care.

Knowledge audits (**see appendix 7.3**) were undertaken across six EDs with a range of expert and novice staff. The aim of this was to examine decision making within the cognitive tasks identified above. Appendix 7.4 provides a fuller summary of the findings.

Figure 6 A simplified cognitive task diagram of ED patients with a suspected pulmonary embolism; those in orange were seen as the most challenging



Initial assessment and selection of investigatory tests

- 4.2.1 Initial assessment of patients arriving at EDs included triage, streaming and some rapid clinical assessment (**see section 1.2**). Rapid clinical assessment included taking a short history, measuring of vital signs and initial tests.
- 4.2.2 Initial assessment was commonly undertaken by nurses with varying levels of experience. Nurses described how assessing patients with non-specific symptoms required expertise to ensure appropriate tests were requested, particularly when there were time pressures.
- 4.2.3 The investigation found that the decision to do a D-dimer test and early suspicion of a PE was influenced by workload, staff experience and ED performance targets. The decision had ramifications for later assessment of patients, influencing the later consideration of PE. Nursing staff rarely found out the outcomes of their decisions or what happened to patients.
- 4.2.4 Expert ED nursing staff described how they tried to focus on symptoms rather than diagnoses, recognising that fixating on a diagnosis too early may influence later decisions. They described that they would only request a D-dimer test after consultation with a senior doctor as they recognised the need for an assessment to be undertaken to determine its need.
- 4.2.5 Novice ED nursing staff told the investigation that they sometimes found the initial assessment of patients difficult. They felt that training had not taught them how to assess symptoms and signs to select investigatory tests; rather they were expected to follow published work procedures. The ACTA subject matter advisor told the investigation that the work procedures in settings such as EDs are unlikely to be able to provide the necessary detail to help staff care for patients with the variety and complexity of conditions that they see in EDs. This meant that training was important.

Further assessment and review of results

- 4.2.6 Further assessment of patients was undertaken after the initial assessment. At this point patients were asked to provide a full description of their symptoms, were examined, and investigatory tests results were reviewed. Further tests were then requested and initial treatment was started if required.
- 4.2.7 Further assessment was undertaken by staff with a range of experience. They included advanced clinical practitioners (ACPs), GPs and doctors of all levels. The ED subject matter advisor told the investigation that this assessment was often undertaken by novice staff, particularly for well patients who may have subtle symptoms and signs.



- 4.2.8 ED staff described that the role of further assessment was to pull together information to identify a main diagnosis. They considered that prolonged experience of working in ED may help the gaining of familiarity with the varied signs and symptoms that patients with PE may show. They also described that expertise was needed to help ensure atypical symptoms and signs, and observations and test results outside of expected ranges, were identified, understood, and acted on appropriately.
- 4.2.9 The investigation was told that the suspicion of, and decision to pursue, a diagnosis of a PE was influenced by workload, years of experience, expertise in the skill of decision making, ED performance targets, available test results and previous decisions about a patient's potential diagnosis.
- 4.2.10 Expert staff described the challenge of building a mental list of potential diagnoses for a patient with non-specific symptoms when information was limited. They likened this to situations that GPs may commonly encounter (**see appendix 7.4** for a knowledge audit undertaken with a GP). They knew they could rarely be 100% sure of a diagnosis, but aimed to refine their mental list using specific questions to account for each symptom, sign and test result. They were concerned with any unexplained finding or finding that did not match with a patient's age, fitness levels or their own reflections on what was wrong with the patient.
- 4.2.11 Experts also used subtle cues to build their understanding of how unwell a patient may be. These cues may be missed by someone with less experience. For example, watching patients walk or talk gave insights into how unwell they were. Some experts also actively sought peer review (opinion from expert colleagues) to check and challenge their diagnoses.
- 4.2.12 For novice staff, the investigation identified several factors that made further assessment difficult (**see appendix 7.4**); a summary is presented here. Novice staff described receiving little training and practice in making decisions. Their education had led them to look for "textbook" presentations. They therefore might reassure themselves that a significant diagnosis is not present in patients who do not "fit the rules". For example, several novices told the investigation that they would not expect a patient who looked well to have a PE or aortic dissection.
- 4.2.13 Novice staff also described how the tests requested at a patient's initial assessment may influence their thinking. For example, the investigation observed that PE had been added to a list of potential diagnoses for a patient after blood test results had been received. The doctor described that they had not initially thought the patient had a PE, but because they had a raised D-dimer test result (ordered at initial assessment), they felt obliged to consider PE as a diagnosis. Alternative causes of the raised D-dimer result were not considered.



4.2.14 Experts told the investigation that they thought novices felt that they had to make a diagnosis to account for a patient's symptoms and test results. This meant that certain tests, such as D-dimer, were sometimes used outside of their intended scope for reassurance. Experts considered that novices may feel uncomfortable exposing their uncertainty about a diagnosis.

4.2.15 The investigation was also told about the influence of COVID-19 on decision making. Early in the pandemic a consultant said: "We have all become novices." The new disease was not part of the experts' experiences and accounting for the new symptoms and signs was therefore challenging. The similarity of PE and COVID-19 symptoms also made identifying PE difficult during the peaks of the pandemic.

Prescribing of interim anticoagulation

4.2.16 The National Institute for Health and Care Excellence (NICE) describes the role of interim anticoagulation for patients with suspected PE and the circumstances in which it should be prescribed and administered (**see 1.1.10**) (National Institute for Health and Care Excellence, 2020a). The decision to prescribe will be made by a doctor or ACP.

4.2.17 Experts described being uncomfortable about prescribing anticoagulation to a patient with an unknown diagnosis. They therefore tried to balance the risk of giving anticoagulation to a patient who ultimately may not need it or to whom it may cause harm, with not giving it to a patient who may need it. The investigation attempted to examine experts' individual strategies to balance risks; nothing formal was described. Some experts used the term "gut feelings" which the investigation considered described their mental shortcuts based on experience.

4.2.18 For novices, some described that prescribing interim anticoagulation was "easy" because NICE told them what to do. The ACTA subject matter advisor suggested that this may be because, unlike experts, they had limited experience of the potential risks. Other novices described the decision to prescribe as challenging; they felt that their training had not equipped them to make judgements about clinical risk.

Summary of findings

4.2.19 The ACTA showed how decisions made in the diagnosis and treatment of PE in ED are influenced by the expertise (knowledge and skill) of decision makers, the available information relating to a patient, and interpretation of and action taken on that information. Decisions were further influenced by the individual circumstances of the staff, training, experience, and wider factors in the healthcare system.

What do experts do?

4.2.20 When presented with familiar situations, experts described knowing what common courses of action to follow. Where situations were less familiar and when there was time, experts described thinking through possibilities in a more analytical way. They were often unable to describe how they structured that analytical thinking.

4.2.21 The terms “gut feeling” and “gestalt” (here meaning an appreciation of the whole problem, rather than its parts) were heard several times. They related to how decision makers built overall perceptions of situations based on factors such as experience, history and examination findings, and other situational prompts. The ACTA subject matter advisor described that a gestalt allows for interpretation of the various sources of information, not just individual symptoms.

4.2.22 The ACTA identified ways experts think and act to support their decision making. These findings may be of value when considering how to develop expert decision-making skills. In summary, experts:

- were alert to the possibility of life-threatening diagnoses and sought to rule these out
- sought to account for unexplained symptoms, signs and results
- were comfortable saying they were unsure of a diagnosis
- recognised the risk of fixating on a diagnosis and compared information critically
- sought opportunities to challenge their decisions with peers
- appreciated when their cognitive abilities may be being affected
- were less worried about making unpopular decisions
- took into account various sources of information including their own knowledge from training, previous experience where applicable, and organisational rules and guidance.

Why are decisions sometimes difficult?

4.2.23 Through the ACTA, novice staff told the investigation of limited recognition of the role of heuristics, limited training in how to make analytical decisions and limited opportunities to develop their decision-making skills in training environments and in clinical practice. These gaps in training and development were further explored by the investigation and findings are described in **section 4.3**.




4.2.24 The ACTA again demonstrated wider factors in the healthcare system that further influenced the decision making of novices and experts. Several of these factors were similar to those identified in the review of the serious incidents, including demand for ED services, capacity in the ED, COVID-19 and standardised processes. These factors were explored further and findings are described in section 4.4.

4.3 Developing decision-making skills

This section describes the findings from the investigation's exploration of how healthcare staff are trained in decision making, and how expert skills are developed. Five factors are explored in turn (see figure 7). While the investigation's focus was on decision making in the diagnosis and treatment of PE in EDs, development of more expert decision-making skills may further support staff when they see patients with other non-specific or atypical symptoms and signs.

Figure 7 Factors examined by the investigation for the development of decision-making skills



Decision making in healthcare curricula

- 4.3.1 The ACTA found that staff had limited familiarity with how people make decisions, and the role of heuristics. Staff also described limited opportunities to practice making decisions. The investigation therefore aimed to understand how the skill of decision making is taught to students and staff.
- 4.3.2 The investigation reviewed several healthcare curricula and standards for training (General Medical Council, 2021; 2018; 2017; Joint Royal Colleges of Physicians Training Board, n.d.; Royal College of Emergency Medicine, 2021a; 2019; Royal College of Nursing, n.d.; Royal College of Physicians, n.d.; 2021). These curricula covered medicine, nursing, advanced clinical practitioners (ACPs – staff with non-medical backgrounds who have skills and knowledge to undertake advanced roles) and physician associates (healthcare professionals who work alongside doctors). They all included decision-making competencies to a lesser or greater degree. For example, the ACP curriculum described knowledge expectations such as to ‘define the steps of diagnostic reasoning’ and ‘demonstrate ability to identify one’s own biases and inconsistencies in clinical reasoning’ (Royal College of Emergency Medicine, 2019).
- 4.3.3 Students and novice doctors told the investigation that their training had not taught them the theory behind how people make decisions, and had not prepared them to make decisions that would affect patients. They described limited appreciation of the differences between the types of decision making or the factors that influence decision making, and had received limited opportunities to practise making decisions.
- 4.3.4 The investigation was told that national curricula and standards were often written by professional bodies. However, implementation and delivery of training is the responsibility of the individual training establishment, for example a university. Professional bodies carry out inspections to assure themselves that establishments are appropriately delivering their curricula.
- 4.3.5 The investigation found a gap between the expectations of curricula, particularly those focused on medical training, around decision-making skills, and how prepared novice doctors felt to make decisions in clinical practice. The investigation identified inconsistencies in how decision making is taught at undergraduate and postgraduate levels. The development of decision-making skills was heard to be via ‘on the job’ experience, implicitly learnt, and with a belief that years of experience equated to expertise.



Simulation and decision making

- 4.3.6 ED leads told the investigation about the use of simulation to train staff in common ED situations. Simulations are 'pretend' versions of healthcare situations that give staff a chance to practise how they would respond. They take place either in the clinical environment or a bespoke simulation suite. Every ED the investigation visited had access to simulation facilities.
- 4.3.7 ED staff told the investigation that they generally welcomed simulation and saw value in it. However, they were rarely able to access it because the need for care provision and workforce shortages often meant simulation and other mandatory training was cancelled. When simulation did take place, it was rarely multi-professional – that is, it only involved some groups of staff. This created unrealistic scenarios because unlike real ED situations, the simulations did not involve the wider multidisciplinary team.
- 4.3.8 Where ED staff had experienced simulation sessions, these focused on emergency teamworking scenarios such as cardiac arrests (when a person's heart stops beating) and trauma. Staff had not experienced any other types of simulation, such as less critical scenarios, those involving patients with non-specific signs and symptoms, or the use of simulation for the evaluation of systems and processes.
- 4.3.9 The research literature provides examples of the positive role of simulation in developing decision-making skills (for example, Murray et al, 2018). Health Education England (HEE) has described the potential benefits of simulation in the development of professional capabilities, multi-professional team working and evaluation of systems and processes. However, HEE has also described unequal access to simulation across the country, varying views on its value, and no agreed approach for how best to implement programmes. In response HEE has developed a vision for simulation in England (Health Education England, 2020).

Serious games

- 4.3.10 HEE told the investigation about other training opportunities available to develop decision-making skills. These included 'serious games', which are games that have been developed for the purpose of something other than entertainment, for example education.
- 4.3.11 Serious games support learning because they are motivating, active, attention grabbing, can be repeated and allow for immediate feedback (Drummond et al, 2017). They are used in several industries; for example, the nuclear industry uses virtual facilities to train inspectors. In healthcare they have been used for patient, public and staff education (Sharifzadeh et al, 2020).



4.3.12 HEE said that serious games could be valuable in helping healthcare novices to develop decision-making skills. This is supported by research that shows the positive effects of serious games on developing decision-making skills (for example, Kaczmarczyk et al, 2016; Crichton et al, 2000). Further positives include limited need for expensive simulation equipment and the option to undertake training as and when the learner wishes (Wang et al, 2016).

Developing decision making in other industries

4.3.13 The ACTA subject matter advisor told the investigation that some other industries deliver training to help accelerate the development of decision-making expertise. The investigation recognises that learning from other industries may not be directly transferrable to healthcare and that any transfer requires evaluation to ensure safety and effectiveness. However, learning from other industries may provide insights and be of value to healthcare.

4.3.14 Research shows that it is possible to help people to accelerate the development of their decision-making skills. For example, studies suggest that 'decision-skills training' (Phillips et al, 2004) and 'accelerating the acquisition of expertise' (Hoffman et al, 2013) can help people to develop recognition-primed decision making skills and more sophisticated heuristics, with a reduced likelihood of them being misapplied.

4.3.15 To understand how industries accelerate skills development, the investigation engaged with military helicopter aviation (case study 1) and the fire service (case study 2). In summary, the investigation found that there were established approaches to help people with limited experience to develop their decision-making skills. The industries developed understanding of decision making, more sophisticated heuristics for recognition-primed decision making, and analytical decision-making skills by:

- Teaching students how people make decisions, the role of heuristics and potential decision-making traps.
- Using mnemonics and 'generic decision tools' to support structured, analytical decision making. These methods and tools are embedded in teaching and practice.
- Including use of the generic decision tools in staff induction and competency assessment. This leads to repetitive practice and the tools are also used to provide feedback by instructors.
- Requiring real-world and simulated application of generic decision tools to maintain competencies, with a minimum yearly simulation requirement. Simulations offer exposure to novel, rare and common situations, enabling staff to recognise them more easily in real life.



4.3.16 The use of simulation in these industries was a critical part of initial and ongoing staff training, and mandatory competency assessments. Staff who did not meet the necessary level of competency would be prevented from undertaking certain responsibilities until the competency was met.

Accelerating decision-making skills in healthcare

4.3.17 The evidence gathered by the investigation suggests that there may be opportunities to support healthcare staff to develop their decision-making skills through:

- the use of simulation to expose staff to situations, for example emergency scenarios, to develop recognition
- the practice of using analytical decision making, for example when considering non-specific or atypical signs and symptoms such as those of PE, where there is time for staff to consider the diagnosis and discuss options with their supervisors.

4.3.18 The learning from the case study industries shows potential value in a comprehensive education focus on understanding decision making, a generic decision tool, and practice in making decisions in a safe environment. The investigation found that while healthcare curricula describe decision-making competencies, novice staff did not feel confident making decisions.

4.3.19 The ACTA subject matter advisor told the investigation that generic decision tools are useful to structure analytical decision making, particularly when the decision maker is confronted with new, unfamiliar situations with time to consider actions. They act as 'scaffolding' for novices.

4.3.20 Various generic decision tools exist outside of healthcare such as DECIDE (define, establish, consider, identify, develop and evaluate) (Guo, 2020), OODA loops (observe, orient, decide and act), and TDODAR (**see figure 8**). The investigation found no established generic decision tool in healthcare and no formal training in using tools to make analytical decisions. Some staff were aware of the tools used by other industries.

Figure 8 TDODAR mnemonic (Civil Aviation Authority, 2014)

- T** – time; do we have time to make a decision or is immediate action needed?
- D** – diagnosis; using available information and challenging any presumptions
- O** – options; weighing up options to address the problem
- D** – decide; choose the most appropriate option
- A** – act; proceed with allocation of tasks to the most appropriate person
- R** – review; assess the outcome, particularly in light of new information

4.3.21 The investigation reviewed the research literature for evidence of impact of generic decision tools such as TDODAR, DECIDE and OODA in industry and healthcare. While the tools are discussed positively by many authors and international bodies, the investigation was not able to find any evaluation studies. In general, tools were felt to support analytical thinking, but potentially required ‘updating’ to bring them in line with modern thinking around decision making (Bryant, 2006). They also required training in their use. Research literature was available for the fire service’s decision control process (**see case study 2**) (Cohen-Hatton and Honey, 2015).

4.3.22 The investigation found that there may be opportunities to improve the use of simulation to support the development of decision-making skills in healthcare. HEE told the investigation that it was keen to enhance the use of simulation and to introduce alternative learning methods to support education in decision making. The investigation recognises the need for staff to have protected time to access training for their development.

4.3.23 The findings led the investigation to make the following safety recommendation and safety observation. The investigation worked with HEE to form the safety recommendation with the aim of supporting staff in EDs to develop their decision-making skills and their recognition of conditions such as PE when faced with non-specific or atypical symptoms and signs. Regarding the safety observation, the investigation recognises the importance of developing an evidence base for the use of a generic decision tool in healthcare to ensure appropriate and safe implementation.

HSIB makes the following safety recommendation

Safety recommendation R/2022/188:

HSIB recommends that Health Education England works with appropriate professional bodies to develop and implement a strategy for supporting the education and training of clinical practitioners that can facilitate the development of decision-making skills. This strategy should consider the use of innovative approaches such as simulation and immersive learning.

HSIB makes the following safety observation

Safety observation O/2022/155:

It may be beneficial for healthcare to learn from other industries and develop its own evidence base on strategies to accelerate the development of expert decision-making skills. These strategies may include:

- development of a generic decision tool for implementation in healthcare training and clinical practice to support analytical decision making
- incorporation into education programmes of theory around how people make decisions and influences on decision making
- the use of simulation as a regular intervention to support practice and development of decision-making skills across scenarios with different levels of complexity
- consideration of the role of simulation in competency assessments for key skills.

Case study 1 – military helicopter aviation

825 Naval Air Squadron (NAS) of the Royal Navy's Fleet Air Arm flies the AgustaWestland Wildcat helicopter. 825 NAS is responsible for aircrew and engineer training on the Wildcat.

To help support how aircrew deal with emergency handling and the making of decisions when faced with any situation, 825 NAS described the use of mnemonics. The mnemonic, **RCSDF** was used to structure the decision thought process and guide the actions of the aircrew.

R – **R**ecognise the emergency; are any immediate actions required?

C – **C**ontain the emergency to prevent further damage

S – achieve ongoing **S**afe flight of the aircraft



D - **D**iagnose the emergency properly

F - use cognitive aids to assist; available through written **F**light reference cards

The investigation observed the use of RCSDF in the helicopter full-motion simulator. Simulation is a core part of initial training and ongoing competency assessments with a minimum annual simulation requirement. In simulations, aircrew are faced with malfunctions to test their decision making during tactical scenarios. They deal with the situations by working through RCSDF step by step. Regular simulator sorties not only expand aircrew experience, but also build 'muscle memory' in dealing with the most serious emergencies where immediate actions are required, such as fires or engine failures.



Picture: Credit (Commanding Officer, 825 Naval Air Squadron, RNAS Yeovilton).


Case study 2 – the fire service

The fire service described to the investigation that decisions are not always made in a linear way (Cohen-Hatton et al, 2015). They rely on various sources of information, received in dynamic and time-pressured situations. In particular, novel and more uncertain situations can result in higher levels of stress which reduce people's capacity to process information and make decisions (Butler et al, 2021).

To support the decision making of incident commanders, the service uses the 'decision control process' (DCP, see figure) (National Fire Chiefs Council, n.d.a). The DCP involves intelligence gathering (situation assessment), planning (based on situation and priorities), mental checking (to ensure the decision is appropriate) and implementation of the plan. It is orientated around three simple questions that incident commanders ask themselves.

The DCP has been evaluated and found to increase appreciation of the situation, result in clearer plans, and reduces the factors that contribute to decisional inertia (hesitancy to make decisions) (Cohen-Hatton and Honey, 2015). Incident commanders are taught the DCP in training and use it in practice via tactical decision games (serious games) and simulated scenarios (National Fire Chiefs Council, n.d.b). Incident commanders are expected to attend a minimum number of incidents per year to maintain the skill; these may be real-world or simulated incidents.

Decision control process, adapted from National Fire Chiefs Council (n.d.a)



Local development of decision-making skills

4.3.24 The ACTA identified further opportunities for individual EDs to support the development of their novices' decision-making skills. These are described below and contributed to the associated safety observation. The ED subject matter advisor described that EDs are ideal learning environments because of the breadth of different symptoms and conditions a patient may arrive with, and this should be taken advantage of for learning. The investigation acknowledges that creating opportunities to improve this type of support for novices requires resource, in particular dedicated time.

Feedback on decisions

4.3.25 The research literature suggests that a fundamental factor in the development of RPD is the ability to review and evaluate the outcome of one's own decisions (Klein, 1997). The investigation observed several ED patients with non-specific symptoms who were diagnosed with suspected PEs. The majority were then referred on or discharged, and the final responsibility for diagnosis passed to someone else.

4.3.26 Some ED staff tracked the outcomes of their patients, generally in their own time. A consultant described to the investigation how they encouraged staff to review cases, including "barndoor" cases (those with a seemingly obvious diagnosis), to see if decisions led to expected outcomes.

4.3.27 The investigation found barriers to seeking information on the outcomes of patients. These included time pressure, motivation at the end of a shift, fear of finding out the outcome, and difficulty accessing medical records. Feedback on performance, including patient outcomes, is critical to the learning process and development of expertise in decision making.

Clinical supervision

4.3.28 Clinical supervision is a process by which professional skills are developed. It includes reflection on practice between supervisors (experts) and practitioners (novices) (Sullivan and Garland, 2013). Development of expertise in decision making requires opportunities for more novice staff to reflect on their practice with experts.

4.3.29 To support supervision and safety, the Royal College of Emergency Medicine has a standard for the 'sign off' of certain ED patient groups (Royal College of Emergency Medicine, 2016). Patients in these groups should be reviewed by a senior doctor (usually a consultant) before being discharged. One of the groups is patients with atraumatic chest pain (that is, chest pain not associated with an injury) aged 30 years and over.



4.3.30 The investigation saw clinical supervision being delivered via ‘shopfloor’ discussions, case reviews and ‘mortality and morbidity’ meetings. The ACTA subject matter advisor described the value of these discussions/meetings to develop RPD. Learning value could be further improved through structuring discussions to focus on the decisions made. In addition, discussions with experts give staff the opportunity to explore subtle aspects of cases that may not be apparent to novices.

4.3.31 In practice the investigation saw supervision discussions managed in different ways, ranging from structured discussions to direct and non-exploratory feedback. The investigation heard that supervisors had rarely received formal education in how to support reflective discussions with novices about their decisions.

Psychological safety

4.3.32 The investigation heard from the fire service about the role of psychological safety in supporting decision making (**see case study 2**). Psychological safety refers to factors that influence how individuals and teams feel they will be perceived when discussing their thoughts (Edmondson, 1999). It is developed through supportive discussions, one-to-one time, formation of relationships, and the valuing of discussions (O’Donovan and McAuliffe, 2020).

4.3.33 Psychological safety was described as necessary for learning, skill development and to support staff to raise concerns in efforts to make healthcare safer. During the investigation’s observations of clinical supervision and case review meetings, psychological safety was perceived to vary. Some situations were observed to be undertaken with non-judgmental questioning of novices, while others were more judgmental. Departmental cultures need to support non-judgmental discussion to encourage staff to seek help, particularly at times of pressure.

HSIB makes the following safety observation

Safety observation O/2022/156:

It may be beneficial if the findings of this investigation are used to support the development of staff expertise in decision making through:


- building understanding of how experts think and make decisions
- supporting reflection on the outcomes of simple and complex decisions
- development of clinical supervision skills of senior staff
- regular multidisciplinary case review.



4.4 Factors in the wider healthcare system that influence decision making

This section describes the findings in relation to factors in the wider healthcare system that support or inhibit diagnostic decision making in EDs. Five factors are explored in turn (see figure 9). These factors emerged from the review of the serious incidents, the ACTA and observations in EDs.

Figure 9 Factors examined by the investigation that influenced decision making



Demand and decision making

4.4.1 The investigation witnessed the significant demand on EDs, which was a result of high patient numbers. ED staff told the investigation that high demand affects the time and information available to them to make decisions. As EDs became busier throughout the day, staff were seen to spend less time with each patient. As a result the thoroughness of assessments was reduced to improve efficiency; this is a recognised risk (Hollnagel, 2009).

4.4.2 Since May 2021 the numbers of patients visiting EDs 'have been the highest on record for their respective month' (Royal College of Emergency Medicine,

2021b). Reports by national bodies describe how demand has exceeded capacity, with inadequate resources, workforce shortages, insufficient hospital beds and limited understanding of local demand contributing to the pressures on EDs (Moulton and Mann, 2021; Royal College of Emergency Medicine, 2020).

- 4.4.3 The examination of ED demand and capacity was outside of the scope of this investigation. However, the investigation recognises that, no matter the expertise of decision makers, degraded decision making may sometimes occur if demand and resource challenges are not addressed.
- 4.4.4 ED staff told the investigation that the national operating standard for EDs (**see 1.2.2**) was another factor that influenced their decisions. Staff described having to make rapid decisions about patients who had been in their EDs for almost 4 hours. The investigation found that staff sometimes felt pressurised to meet the standard and described a sometimes punitive response if it was not met.
- 4.4.5 The national operating standard in England is contentious. There is evidence that it has benefited patients by reducing long waits and overcrowding, which are known to be associated with harm. However, the standard has not been consistently met for some time (NHS England and NHS Improvement, n.d.) and may have led to a focus on performance, rather than quality of care and good outcomes for patients (Moulton and Mann, 2021). At the time of writing there are proposals for the standard to be replaced (NHS England and NHS Improvement, 2020; O'Dowd, 2020) and new operational priorities are planned (NHS England and NHS Improvement, 2022).

National guidance and decision making

- 4.4.6 The National Institute for Health and Care Excellence (NICE) provides guidance on the diagnosis and treatment of adults with PE (and deep vein thrombosis) (National Institute for Health and Care Excellence, 2020a). The investigation found that ED staff were familiar with the guidance, but rarely applied the decision-making scores and criteria described in it.

Scores and criteria supporting decisions

- 4.4.7 Where there is a low suspicion of PE and another diagnosis is more likely, NICE recommends the use of the pulmonary embolism rule-out criteria (PERC) to help determine whether further tests for PE are needed (National Institute for Health and Care Excellence, 2020a, 2020b). NICE states that PERC should be considered where 'the clinician estimates the likelihood of PE to be less than 15% based on the overall clinical impression, and other diagnoses are feasible'. ED staff described concerns about the guidance for PERC that limited their use of it to inform their decisions:

- In practice determining a 'less than 15%' likelihood of PE in a patient is challenging. NICE does not provide further guidance on this, but does provide the background to the recommendation (National Institute for Health and Care Excellence, 2020b).
- The criteria for when PERC can and cannot be used is unclear. ED staff described uncertainty about situations such as patients with active cancer or who are on the oral contraceptive pill.

4.4.8 The ED subject matter advisor told the investigation that they had seen PERC support clinical decision making and promote evidence-based care. However, the ED and haematology subject matter advisors said that it can be inappropriately applied where staff do not understand its role and there is limited clarity around its use. The investigation also noted that even when correctly applied, there will still be a small proportion of patients who have a PE but meet the conditions of PERC (Kline et al, 2004).

4.4.9 Where there is a suspicion of PE, NICE recommends the use of the Wells score (**see 1.1.6**) to guide whether a D-dimer test may be used (National Institute for Health and Care Excellence, 2020a). During its observations, the investigation did not see staff documenting a Wells score before requesting a D-dimer test and did not hear it being used to support diagnostic decisions for PE.

4.4.10 ED staff described concerns about the validity of the Wells score that limited their use of it. Because it is over 20 years old (Wells et al, 2001), staff questioned whether it accounted for the types of patients now commonly seen in EDs such as those with obesity, on the oral contraceptive pill, with suspected but not confirmed cancer, and with reasons for not developing a high heart rate.

4.4.11 Several consultant doctors told the investigation of other scoring systems that they perceived as more valid than the Wells score. The haematology subject matter advisor and another national expert in thrombosis told the investigation that the Wells score was still the most appropriate to use.

Decision to start patients on interim anticoagulation

4.4.12 The ACTA identified that the decision to give a patient interim anticoagulation was sometimes challenging for ED staff. Once a PE has been suspected, NICE recommends anticoagulation if a scan cannot be done immediately (if a patient is likely to have a PE), or if a D-dimer test result is not available within 4 hours (if a patient is unlikely to have a PE) (National Institute for Health and Care Excellence, 2020a). Regarding imaging, the guidance does not define a timeframe for 'immediately'; it may be that a computerised tomography pulmonary angiography (CTPA) is ordered immediately, but that limited capacity means a delay.

- 4.4.13 ED doctors and ACPs were familiar with NICE's recommendation for anticoagulation pending CTPA, but shared occasions where they had withheld anticoagulation. They described their concerns about the safety of a "blood thinning medicine" where a diagnosis is unknown. They felt that NICE's recommendation did not consider the risk to patients of taking anticoagulation if their diagnosis turned out to be something other than PE. In addition, the recommendation did not state the circumstances in which anticoagulation should not be administered.
- 4.4.14 The haematology subject matter advisor stated that the risk from a single dose of low molecular weight heparin (commonly used for interim anticoagulation) was small (Chai-Adisaksopha et al, 2014; Agnelli et al, 2013; Büller et al, 2012) unless there were obvious risk factors, for example major recent surgery. A comprehensive history was needed to identify bleeding risk and other diagnoses. However, it was acknowledged that a comprehensive history requires time which may be limited in an ED.
- 4.4.15 In relation to the administration of anticoagulation pending a D-dimer result, ED staff were often unaware of this NICE recommendation and suspected that it was aimed at primary care. NICE told the investigation that the guidance was for all providers of primary, secondary and tertiary care (for example, GPs, hospitals and specialist care centres) (National Institute for Health and Care Excellence, 2020a).

The influence of COVID-19 on decision making

- 4.4.16 ED staff shared concerns that the emergence of COVID-19 may invalidate some of the recommendations in NICE guidance for assessing the risk of PE. This had the potential to impact on their decisions when diagnosing PE. With the risk of blood clots associated with COVID-19, staff questioned whether D-dimer result thresholds and the Wells score were still appropriate.
- 4.4.17 NICE and the subject matter advisors told the investigation that there was limited research into the validity of scoring systems and D-dimer tests in COVID-19. There are differing opinions about assessment of the risk of blood clots in patients with the virus (Patel et al, 2021) and various societies have published advice (for example, British Society of Thoracic Imaging, 2021).

Summary

- 4.4.18 NICE told the investigation that its document provides guidance based on consideration of available evidence, but that clinicians must make decisions appropriate to the individual circumstances they are presented with. However,

NICE welcomes opportunities to clarify and make its guidance more usable. The investigation found that clarity on some aspects of the NICE guidance would be welcomed to support ED staff in diagnosing and treating suspected PE. As a result, HSIB makes the following safety recommendation with the intent that it clarifies guidance where possible to support decision making, and identifies whether more recent evidence is available to inform guidance updates.

HSIB makes the following safety recommendation

Safety recommendation R/2022/189:

HSIB recommends that the National Institute for Health and Care Excellence reviews the findings of this investigation in relation to its guidance NG158, 'Venous thromboembolic diseases: diagnosis, management and thrombophilia testing', and updates the guidance if required.

4.4.19 The NICE guidance collates national evidence in order to guide the decisions and actions of healthcare staff in practice. However, the investigation identified that the implementation of the guidance in practice was not consistent. This was in part due to the need to clarify some aspects of the guidance for staff, as mentioned above. Other local barriers to implementation of guidance were also identified. These included local guidance that was not in line with national guidance, unpredictable delays to imaging or blood results, and limited local efforts to embed the national guidance. These barriers were and will be location specific; there is therefore value in identifying the barriers in each organisation.

HSIB makes the following safety observation

Safety observation O/2022/157:

It may be beneficial for individual organisations to understand the extent to which national guidance on the diagnosis and management of pulmonary embolism is implemented across their organisations. This would help to identify local barriers to implementation to address. In particular it may be helpful to consider, in line with the findings of this investigation, local engagement with the scoring systems available to help predict the likelihood of a pulmonary embolism.

Design of emergency department processes

4.4.20 The investigation found that the availability, organisation and design of processes in EDs influenced the decisions made by staff. This was identified in relation to processes for the initial assessment of patients, pathways for the outpatient care of well patients with a suspected PE, and handover of patient care.

Initial assessment design and D-dimer decisions

- 4.4.21 The investigation observed initial assessment processes. The investigation regularly heard that the role of initial assessment was to identify the patient's symptoms and request tests relevant to those symptoms, not to make a diagnosis. The ACTA identified that the "labelling" of patients with a diagnosis, such as a suspected PE, and requesting (or not) a D-dimer test later influenced diagnostic decisions. The investigation found inconsistencies in the initial assessment processes.
- 4.4.22 The initial assessments were found to commonly merge triage, streaming and rapid initial assessment; the Royal College of Emergency Medicine (RCEM) describes these as distinct processes (**see section 1.2**). For patients with chest pain and/or shortness of breath this resulted in variation in where patients were seen (for example major or minor areas), who they were seen by, and the tests requested.
- 4.4.23 The investigation observed variation in how initial assessment staff requested D-dimer and troponin (protein in the blood measured to help diagnosis of a heart attack) blood tests. There was limited opportunity for staff to explore a patient's symptoms and decide whether a test was appropriate. Staff described the need to do all the tests at once to reduce the risk of "breaching" the national operating standard and to not miss anything.
- 4.4.24 NICE recommends a D-dimer test if a Wells score suggests this is necessary (National Institute for Health and Care Excellence, 2020a). The investigation found no Wells scores completed before D-dimers being requested at initial assessment. Staff were predominantly nurses who said they had had no training in the use of the Wells score. ED doctors told the investigation that a Wells score requires clinical judgement and cannot be done without training and time.
- 4.4.25 The investigation found that the absence of a Wells score at initial assessment meant D-dimers were requested without formal consideration of their need in line with NICE guidance. The observed outcomes of this included:
- staff at further assessment feeling it necessary to "rule-out" a PE where a D-dimer level was raised
 - application of a rule-based decision (raised D-dimer means PE) without consideration of other potential diagnoses and causes
 - anxiety if an elevated D-dimer level was later "ignored" if it was felt not to be a PE.

- 4.4.26 To address the issue of requesting D-dimers without consideration of their need, some EDs collected specific blood samples that could be tested for D-dimer levels later if felt to be required. It was heard that this increased costs and demand on laboratories.
- 4.4.27 The investigation found workforce shortages across EDs and noted recruitment and retention challenges (NHS Digital, n.d.). ED leads told the investigation that this resulted in less experienced nursing staff undertaking initial assessment. Senior ED staff felt that these staff struggled to assess risk and therefore requested a “battery” of tests without considering why.
- 4.4.28 The investigation sought to understand how nurses were trained and supported in their initial assessment roles. Nurses undergo local training with the focus on triage. Variation was seen in the content of training across EDs and there was limited inclusion of competencies related to focused assessment and test selection. Staff were expected to access and follow work procedures to know what to do.
- 4.4.29 The design of and access to work procedures, such as standard operating procedures and guidelines, is considered further in sections 4.4.52 to 4.4.55. The investigation found work procedures specific to the diagnosis and treatment of PE in all EDs. The procedures’ specific focus on a diagnosis (PE) was noted to be at variance with the expectation of initial assessment to focus on symptoms and to not make a diagnosis. The investigation considered that the focus of the procedures may further prime staff to think PE, or not, at initial assessment.
- 4.4.30 The investigation had the opportunity to observe formal rapid clinical assessments at several EDs. Rapid clinical assessment was a limited resource and was led by a doctor (registrar or consultant level) or ACP. Staff described the benefits of rapid assessment, including focused assessments and rational requesting of tests. The ED subject matter advisor agreed with the value of having expert decision makers at initial assessment and that this is the ambition of the RCEM.
- 4.4.31 The investigation shared its findings with the RCEM and heard that there is limited evidence around what is the most effective model for initial assessment (Royal College of Emergency Medicine, 2017). The challenges with requesting and interpreting D-dimer results were further recognised by many of the staff, experts and stakeholders who engaged with the investigation. RCEM described its various workstreams and resources that may help address the identified issues:
- a service design and configuration committee, the function of which is to look at such service issues

- the Consultant Sign-Off Standard which includes atraumatic chest pain in those over 30 years (**see 4.3.29**) (Royal College of Emergency Medicine, 2016)
- a learning module on PE for staff in EDs which includes aspects related to D-dimer testing (Royal College of Emergency Medicine, 2019).

4.4.32 As a result of the findings, HSIB makes the following safety recommendation. Because of the scope of the investigation, the safety recommendation focuses on PE. However, it also has the potential to inform future standardisation of initial assessment processes, based on clearer evidence, for any condition with non-specific symptoms and signs where tests are needed to help narrow the diagnosis.

HSIB makes the following safety recommendation

Safety recommendation R/2022/190:

HSIB recommends that the Royal College of Emergency Medicine promotes best practice around diagnostic decision making with respect to patients with potential symptoms and signs of pulmonary embolism.

Outpatient PE pathways

- 4.4.33 Across the EDs the investigation saw patients with suspected PEs being discharged for outpatient diagnosis and treatment. In these situations, patients received anticoagulation and returned for imaging in the next 24 to 72 hours. This is in line with NICE (National Institute for Health and Care Excellence, 2020a, 2020b) and British Thoracic Society (Howard et al, 2018) guidance.
- 4.4.34 The NHS's strategy is to enable more patients to be treated outside of hospitals; the benefits of this include fewer hospital admissions and improved patient experience. Same-day emergency care services, where patients with relevant conditions can be rapidly assessed, treated and discharged, are part of the NHS's Long Term Plan (NHS England and NHS Improvement, 2019) and recommended by the 'Getting It Right First Time' programme's national specialty report into emergency medicine (Moulton and Mann, 2021).
- 4.4.35 ED staff described their positive perceptions of outpatient/same-day care for patients with suspected "barndoor [obvious] PEs." However, they also expressed a perceived pressure to discharge patients that sometimes influenced their decisions. They recognised the risks of discharging patients on anticoagulation medicines without a confirmed diagnosis of PE. Staff challenged the risk tools used to support the discharge decisions, such as the 'simplified pulmonary embolism severity index (sPESI) (Howard et al, 2018)', over concerns that they were based on the risk of death in patients with a



confirmed diagnosis of PE, not where PE was suspected and an alternative diagnosis could be present. Leads from NHS England and NHS Improvement's programme for same-day emergency care told the investigation of the need for local services to ensure they have safe discharge protocols in place for use of same-day emergency care services where diagnoses cannot be confirmed on the same day in the ED.

4.4.36 The investigation found a preference for discharging patients with suspected PE from EDs for outpatient diagnosis, including where ED staff perceived that there was local capacity for same-day imaging. Some of these patients then went on to wait for 72 hours for a scan; this created a safety risk with patients on anticoagulation waiting for a diagnosis. As per NICE guidance, patients with a suspected PE should receive an immediate CTPA. As a result, HSIB makes the following safety observation, recognising that various factors influence imaging availability.

HSIB makes the following safety observation

Safety observation O/2022/158:

It may be beneficial for emergency departments and same-day emergency care units to have rapid access to recommended imaging for patients who require it for the diagnosis of pulmonary embolism.

Handover of patient care

4.4.37 Handovers of care were observed across the EDs, most commonly between staff at the change of shift. The investigation was told of examples where loss of information at handover had influenced later decisions and resulted in delayed or missed diagnoses of PE. This was demonstrated by Martin's case, which is detailed in **section 3**.

4.4.38 ED staff described mixed practices of written and verbal handover. The investigation found attempts in various EDs to improve task handover, such as the introduction of 'huddles' (group discussions), electronic task allocation and digital systems for test ordering that required confirmation that test results had been seen and that further action had been taken accordingly.

4.4.39 The addition of technology was thought to be beneficial in the support of handover. However, technology alone will not always mean handover is effective in the support of the actions and decisions of staff. The ACTA subject matter advisor stated that the critical elements of a handover are the people involved, and ensuring 'closed-loop communication' means that the right person has received the right information, has understood it, and is able to carry out the task with appropriate training and resources.



Decisions to request imaging

- 4.4.40 The serious incident reports and interviews with ED staff identified occasions where ED doctors and ACPs had faced barriers to accessing CTPAs for patients who they suspected had a PE. Staff described that these barriers may have influenced their decisions to diagnose a patient with a suspected PE.
- 4.4.41 ED staff described times when they had felt a CTPA was required, but radiology staff had challenged the need. Those challenges related to the timing of the CTPA (immediate versus a later outpatient scan) and where radiology felt a CTPA was not justified. ED staff described frustration when challenged as they had undertaken the assessments, were under time pressure and feared missing a PE. Increased requesting of CTPAs because of fear of missing a PE is described in the research literature (Rohacek et al, 2012).
- 4.4.42 The investigation observed CTPAs being undertaken in several hospitals. Radiology staff described that requests for CTPA were often made without use of the recommended decision tools, such as Wells score; this is widely reported (Kauppi et al, 2021; Perera et al, 2017). Radiology staff also wanted to ensure patients were not exposed to unnecessary radiation.
- 4.4.43 The ED subject matter advisor told the investigation that if a CTPA is found to be needed, following use of the appropriate decision tools, then it should be done, ideally immediately. The investigation saw examples of where multiple specialties, including ED and radiology, had worked collaboratively to develop clear expectations of the information required to justify a CTPA. This was thought to support ED staff decision making, improve requesting practices and reduce conflicts.

Access to CTPA

- 4.4.44 The investigation found that a significant factor that made it difficult for ED staff to access CTPA was the available capacity in some hospitals. Limited CTPA capacity was commonly described by ED staff; this was confirmed by radiology staff. The investigation engaged with the Royal College of Radiologists (RCR) and heard of significant shortages of imaging capacity, but that the aim should be to image all patients within 24 hours.
- 4.4.45 An 'Independent review of diagnostic services for NHS England' (Richards, 2020) noted significant shortages in imaging capacity in England, with the country having the lowest CT scanner provision amongst 23 countries. The review recommended a 100% expansion of CT capacity and major expansions in workforce.



4.4.46 Further exploration of imaging capacity and workforce was out of scope of this investigation and there is little value in repeating recommendations already made by Richards (2020). However, the investigation heard that limited CTPA capacity potentially affected ED staff members' decisions to diagnose patients with a suspected PE and request CTPA. Shortages of CT were also identified in HSIB's investigation report, 'Missed detection of lung cancer on chest X-rays of patients being seen in primary care' (**Healthcare Safety Investigation Branch, 2021c**).

4.4.47 In some EDs staff described "no problem" getting a same-day CTPA even for well patients, despite increased demand because of COVID-19. These hospitals generally had CT scanners in their EDs, but despite the perceived capacity, the investigation noted that some staff still used the outpatient PE pathway. The investigation was told this was because staff were unaware of imaging capacity, outpatient diagnosis was the trust's preferred pathway, and because they did not want to have potentially challenging discussions with radiology staff.

National CTPA standard

4.4.48 Radiology staff in hospitals told the investigation about the increasing demand for CTPA, but with fewer scans showing presence of a PE (Dobler, 2019; Perera et al, 2017). In response, the RCR has developed a standard that at least 15% of CTPAs should show presence (positivity) of a PE (Royal College of Radiologists, 2021). The RCR also provides iRefer, under revision at the time of writing, which provides guidance on the requesting of radiological examinations included CTPA for PE (Royal College of Radiologists, n.d.).

4.4.49 The RCR told the investigation that the positivity standard was introduced to help ensure that appropriate patients (following determination of PE risk in line with national guidance) receive imaging and that radiologists must justify exposure of patients to radiation. The standard is also intended to ensure under-requesting of CTPA is identified and addressed. Where local CTPA positivity results are below or significantly above the standard, this should prompt local discussions to refine guidelines for suspected PE.

4.4.50 The RCR stated that they felt the standard was reasonable and was based on a review of evidence of the varying positivity rates across the research literature (for example, Costa et al, 2014). No national data on positivity rates was available, but the RCR has provided an audit plan for the assessment of local practice (Royal College of Radiologists, 2021).



4.4.51 ED doctors told the investigation that they felt that the standard was seen as a “target” and created doubt in some of their minds when deciding whether a CTPA was required. There were concerns that it had been instituted without supporting evidence and consideration of the potential impact on ED decision making.

HSIB makes the following safety observation

Safety observation O/2022/159:

It may be beneficial for the positivity standard for computerised tomography pulmonary angiography (CTPA) (that at least 15% of CTPAs should show a pulmonary embolism) to be evaluated to understand its effects on emergency department decision making.

Work environment and decision making

Work procedures

4.4.52 The investigation found work procedures in all EDs that provided information to staff on how to diagnose and treat PE. Work procedures are a ‘logical step-by-step way of doing things at work, often in the form of written instructions, checklists, decision aids, diagrams, or flow charts’ (Chartered Institute of Ergonomics and Human Factors, 2020). They standardise tasks, enhance best practice and reduce reliance on memory.

4.4.53 Work procedures were found in paper and electronic formats, but the investigation did not observe staff accessing them. The investigation was told by the subject matter advisors and ED staff that this likely contributed to inconsistencies in the assessment of patients and requesting of tests.

ED staff told the investigation that they did not use work procedures because:


- they did not feel they needed to
- they could recall from memory what needed to be done
- there was not enough time at initial assessment to go through them
- the procedures did not address the reality they were facing
- the procedures were difficult to find and not easy to use.

4.4.54 The investigation found that several procedures did not match the realities of the work and were presented in a way that were difficult to follow. The investigation also identified errors in some procedures.



4.4.55 The research literature describes the benefits of well-designed procedures to support tasks, including in relation to PE (Ehrman et al, 2021; Richardson et al, 2020). The Chartered Institute of Ergonomics and Human Factors has published guidance to support the design of effective and usable work procedures (and guidelines) (see figure 10) (Chartered Institute of Ergonomics and Human Factors, 2020). The investigation found that these principles had not been applied to the procedures it had reviewed. As a result, HSIB makes the following safety observation, acknowledging that organisations need to create the conditions within which the work procedures can easily be applied, and that staff need protected time and training to develop effective procedures.

Figure 10 Guiding principles for effective and usable work procedures adapted from the Chartered Institute of Ergonomics and Human Factors (2020)



HSIB makes the following safety observation

Safety observation O/2022/160:

It may be beneficial for healthcare work procedures to be written in line with the principles for effectiveness and usability provided by the Chartered Institute of Ergonomics and Human Factors.

Workspace location and design

- 4.4.56 Observations and discussions with staff demonstrated how the workspace location and the physical environment could support or inhibit decision making.
- 4.4.57 The investigation spent time in areas designated for patients who were not acutely unwell. The investigation was told that a patient being in these locations can influence clinical decisions about their care. For example, well patients in 'minors' or ambulatory areas of ED were less likely to have blood tests. The staff carrying out assessments were therefore more likely to think that patients had less severe conditions. Similarly, the investigation was told that if a patient was in a COVID-19 area, COVID-19 always became a top diagnosis.
- 4.4.58 Physical workspace design was noted to potentially influence decisions. Several EDs had limited quiet areas where staff could retire to consider diagnoses and plans. The investigation observed staff using such areas where they existed and witnessed the role they played in creating a space for informal conversations about patient symptoms and diagnoses.
- 4.4.59 From a human factors perspective, the design of physical environments can support or inhibit staff in the tasks they are undertaking. Environments may support decision making by encouraging informal, opportunistic exchanges of information, and by minimising distractions and interruptions. The investigation acknowledges that, for many reasons, it is difficult to adapt current physical workspaces and environments to better support staff performance.



5 Summary of findings, safety recommendations and safety observations

5.1 Findings

- Recognising that a person may have a PE is challenging, particularly for less experienced staff and when the person's signs and symptoms are non-specific or atypical.
- Deciding whether to initiate treatment for a suspected PE requires a decision that balances risks, and this decision can benefit from expert knowledge and skill.
- Despite expertise and the available tools to help identify patients who may have a PE, a small number of PEs may always be missed.
- Experts use different thought processes and show different behaviours when making decisions compared to more novice staff.
- Decision-making skills in healthcare are commonly developed through experience, without formal training or opportunities to practise making decisions.
- Simulation-based learning has the potential to help staff acquire decision-making skills more quickly.
- Other industry sectors, such as aviation and the fire service, aim to accelerate the development of decision-making skills through structured training and the use of 'generic decision tools' for analytical decisions.
- EDs do not always provide the conditions which support the development of decision-making skills.
- Decision making in EDs is affected by workload, workforce availability, and performance targets.
- ED staff asked for further guidance to be provided on the use of decision aids to support the diagnosis of PE.
- The design of ED processes influences the decisions staff make. There is no standard model of initial patient assessment in EDs; this contributes to variation in the requesting of tests which can affect later decisions.

- Pathways for the diagnosis and treatment of PE in outpatient settings may create a safety risk where patients are discharged on anticoagulation medicines without a confirmed diagnosis; the capacity of imaging services is a significant contributor to this.
- Loss of clinical information when a patient's care is handed over was identified as a further safety risk. This can contribute to harm if tests, such as D-dimer (a blood test used as part of the assessment of likelihood of PE), are not followed up.
- Work procedures for the diagnosis and treatment of PE are not routinely designed in line with human factors principles to support their access and use.
- The physical design of environments may also affect decision making.

5.2 HSIB makes the following safety recommendations

HSIB makes the following safety recommendations

Safety recommendation R/2022/188:

HSIB recommends that Health Education England works with appropriate professional bodies to develop and implement a strategy for supporting the education and training of clinical practitioners that can facilitate the development of decision-making skills. This strategy should consider the use of innovative approaches such as simulation and immersive learning.

Safety recommendation R/2022/189:

HSIB recommends that the National Institute for Health and Care Excellence reviews the findings of this investigation in relation to its guidance NG158, 'Venous thromboembolic diseases: diagnosis, management and thrombophilia testing', and updates the guidance if required.

Safety recommendation R/2022/190:

HSIB recommends that the Royal College of Emergency Medicine promotes best practice around diagnostic decision making with respect to patients with potential symptoms and signs of pulmonary embolism.



HSIB makes the following safety observations

Safety observation O/2022/155:

It may be beneficial for healthcare to learn from other industries and develop its own evidence base on strategies to accelerate the development of expert decision-making skills. These strategies may include:

- development of a generic decision tool for implementation in healthcare training and clinical practice to support analytical decision making
- incorporation into education programmes of theory around how people make decisions and influences on decision making
- the use of simulation as a regular intervention to support practice and development of decision-making skills across scenarios with different levels of complexity
- consideration of the role of simulation in competency assessments for key skills.

Safety observation O/2022/156:

It may be beneficial if the findings of this investigation are used to support the development of staff expertise in decision making through:

- building understanding of how experts think and make decisions
- supporting reflection on the outcomes of simple and complex decisions
- development of clinical supervision skills of senior staff
- regular multidisciplinary case review.

Safety observation O/2022/157:

It may be beneficial for individual organisations to understand the extent to which national guidance on the diagnosis and management of pulmonary embolism is implemented across their organisations. This would help to identify local barriers to implementation to address. In particular it may be helpful to consider, in line with the findings of this investigation, local engagement with the scoring systems available to help predict the likelihood of a pulmonary embolism.



Safety observation O/2022/158:

It may be beneficial for emergency departments and same-day emergency care units to have rapid access to recommended imaging for patients who require it for the diagnosis of pulmonary embolism.

Safety observation O/2022/159:

It may be beneficial for the positivity standard for computerised tomography pulmonary angiography (CTPA) (that at least 15% of CTPAs should show a pulmonary embolism) to be evaluated to understand its effects on emergency department decision making.

Safety observation O/2022/160:

It may be beneficial for healthcare work procedures to be written in line with the principles for effectiveness and usability provided by the Chartered Institute of Ergonomics and Human Factors.



6 References

Agnelli, G., Buller, H. R., Cohen, A., Curto, M., Gallus, A. S., Johnson, M., Masiukiewicz, U., Pak, R., Thompson, J., Raskob, G. E. and Weitz, J. I. (2013) Oral apixaban for the treatment of acute venous thromboembolism. *The New England Journal of Medicine*, 369 (9), 799-808.

Barco, S., Mahmoudpour, S. H., Valerio, L., Klok, F. A., Münzel, T., Middeldorp, S., Ageno, W., Cohen, A. T., Hunt, B. J. and Konstantinides, S. V. (2020) Trends in mortality related to pulmonary embolism in the European Region, 2000-15: analysis of vital registration data from the WHO Mortality Database. *The Lancet Respiratory Medicine*, 8 (3), 277-287.

BMJ Best Practice. (2021) Pulmonary embolism [Online]. Available at <https://bestpractice.bmj.com/topics/en-gb/3000115/diagnosis-recommendations> (Accessed 13 December 2021).

British Society of Thoracic Imaging. (2021) British Society of Thoracic Imaging: Rationale for CTPA in Covid-19 patients [Online]. Available at https://www.bsti.org.uk/media/resources/files/Rationale_for_CTPA_in_Covid_considerations_F.pdf (Accessed 17 August 2021).

Bryant, D. J. (2006) Rethinking OODA: toward a modern cognitive framework of command decision making. *Military Psychology*, 18 (3), 183-206.

Büller, H. R., Prins, M. H., Lensin, A. W. A., Decousus, H., Jacobson, B. F., Minar, E., Chlumsky, J., Verhamme, P., Wells, P., Agnelli, G., Cohen, A., Berkowitz, S. D., Bounameaux, H., Davidson, B. L., Misselwitz, F., Gallus, A. S., Raskob, G. E., Schellong, S. and Segers, A. (2012) Oral rivaroxaban for the treatment of symptomatic pulmonary embolism. *The New England Journal of Medicine*, 366 (14), 1287-1297.

Butler, P. C., Bowers, A., Smith, A. P., Cohen-Hatton, S. R. and Honey, R. C. (2021) Decision making within and outside standard operating procedures: paradoxical use of operational discretion in firefighters. *Human Factors* [Online]. DOI: 10.1177/00187208211041860.

Carayon, P., Schoofs Hundt, A., Karsh, B.-T., Gurses, A. P., Alvarado, C. J., Smith, M. and Flatley Brennan, P. (2006) Work system design for patient safety: the SEIPS model. *Quality & Safety in Health Care*, 15 (Suppl 1), i50-8.

Chai-Adisaksopha, C., Crowther, M., Isayama, T. and Lim, W. (2014) The impact of bleeding complications in patients receiving target-specific oral anticoagulants: a systematic review and meta-analysis. *Blood*, 124 (15), 2450-2458.



Chartered Institute of Ergonomics and Human Factors. (2020) Guidance to help design effective and usable work procedures for health and social care teams [Online]. Available at <https://ergonomics.org.uk/resource/guidance-on-design-of-effective-work-procedures.html> (Accessed 22 February 2022).

Choong, C. K., Calvert, P. A., Falter, F., Mathur, R., Appleton, D., Wells, F. C., Schofield, P. M. and Crawford, R. (2008) Life-threatening impending paradoxical embolus caught “red-handed”: successful management by multidisciplinary team approach. *The Journal of Thoracic and Cardiovascular Surgery*, 136 (2), 527-528.

Clarity Informatics Ltd. (2020) Pulmonary embolism: clinical knowledge summary commissioned and funded by National Institute for Health and Care Excellence [Online]. Available at <https://cks.nice.org.uk/topics/pulmonary-embolism/> (Accessed 24 February 2020).

Clinical Human Factors Group. (2018) Human factors in healthcare: common terms [Online]. Available at <http://s753619566.websitehome.co.uk/wp-content/uploads/2018/06/chfg-human-factors-common-terms.pdf> (Accessed 12 March 2021).

Cohen-Hatton, S. R. and Honey, R. C. (2015) Goal-oriented training affects decision-making processes in virtual and simulated fire and rescue environments. *Journal of Experimental Psychology*, 21 (4), 395-406.

Cohen-Hatton, S. R., Butler, P. C. and Honey, R. C. (2015) An investigation of operational decision making in situ: incident command in the UK Fire and Rescue Service. *Human Factors*, 57 (5), 793-804.

Costa, A. F., Basseri, H., Sheikh, A., Stiell, I. and Dennie, C. (2014) The yield of CT pulmonary angiograms to exclude acute pulmonary embolism. *Emergency radiology*, 21 (2), 133-141.

Civil Aviation Authority. (2014) Flight-crew human factors handbook CAP 737 [Online]. Available at <https://publicapps.caa.co.uk/docs/33/CAP%20737%20DEC16.pdf> (Accessed 6 January 2022).

Crichton, M. T., Flin, R. and Rattray, W. A. R. (2000) Training decision makers – tactical decision games. *Journal of Contingencies and Crisis Management*, 8 (4), 208-217.

Dobler, C. C. (2019) Overdiagnosis of pulmonary embolism: definition, causes and implications. *Breathe*, 15 (1), 46-53.

Dreyfus, S. E. (2004) The five-stage model of adult skill acquisition. *Bulletin of Science Technology & Society*, 24 (3), 177-181.

Drummond, D., Hadchouel, A. and Tesnière, A. (2017) Serious games for health: three steps forwards. *Advances in Simulation*, 2 (3).

Edmondson, A. (1999) Psychological safety and learning behavior in work teams. *Administrative Science Quarterly*, 44 (2), 350.

Ehrman, R. R., Malik, A. N., Smith, R. K., Kalarikkal, Z., Huang, A., King, R. M., Green, R. D., O'Neil, B. J. and Sherwin, R. L. (2021) Serial use of existing clinical decisions aids can reduce computed tomography pulmonary angiography for pulmonary embolism. *Internal and Emergency Medicine*, 16 (8), 2251-2259.

General Medical Council. (2017) Generic professional capabilities framework [Online]. Available at <https://www.gmc-uk.org/education/standards-guidance-and-curricula/standards-and-outcomes/generic-professional-capabilities-framework> (Accessed 18 June 2021).

General Medical Council. (2018) Outcomes for graduates 2018 [Online]. Available at https://www.gmc-uk.org/-/media/documents/outcomes-for-graduates-2020_pdf-84622587.pdf?la=en&hash=35E569DEB208E71D666BA91CE58E5337CD569945 (Accessed 18 June 2021).

General Medical Council. (2021) MLA content map: Medical Licensing Assessment [Online]. Available at https://www.gmc-uk.org/-/media/documents/mla-content-map-_pdf-85707770.pdf (Accessed 8 July 2021).

Guo, K. L. (2020) DECIDE: a decision-making model for more effective decision making by health care managers. *The Health Care Manager*, 39 (3), 133-141.

Health Education England. (2020) Enhancing education, clinical practice and staff wellbeing. A national vision for the role of simulation and immersive learning technologies in health and care: Technology Enhanced Learning (TEL) [Online]. Available at <https://www.hee.nhs.uk/sites/default/files/documents/National%20Strategic%20Vision%20of%20Sim%20in%20Health%20and%20Care.pdf> (Accessed 10 June 2021).

Healthcare Safety Investigation Branch. (2020a) Management of venous thromboembolism risk in patients following thrombolysis for an acute stroke [Online]. Available at <https://www.hsib.org.uk/investigations-and-reports/management-of-venous-thromboembolism-risk-in-patients-following-thrombolysis-for-an-acute-stroke/management-of-venous-thromboembolism-risk-in-patients-following-thrombolysis-for-an-acute-stroke/> (Accessed 17 September 2021).

Healthcare Safety Investigation Branch. (2020b) Delayed recognition of acute aortic dissection [Online]. Available at <https://www.hsib.org.uk/investigations-and-reports/delayed-recognition-of-acute-aortic-dissection/delayed-recognition-of-acute-aortic-dissection/> (Accessed 18 June 2021).

Healthcare Safety Investigation Branch. (2021a) Support for staff following patient safety incidents [Online]. Available at <https://www.hsib.org.uk/investigations-and-reports/support-for-staff-following-patient-safety-incidents/national-learning-report-support-for-staff-following-patient-safety-incidents/> (Accessed 24 February 2021).

Healthcare Safety Investigation Branch. (2021b) Never Events: analysis of HSIB's national investigations, Healthcare Safety Investigation Branch [Online]. Available at <https://www.hsib.org.uk/investigations-and-reports/never-events-analysis-of-hsibs-national-investigations/> (Accessed 24 February 2021).

Healthcare Safety Investigation Branch. (2021c) Missed detection of lung cancer on chest X-rays of patients being seen in primary care [Online]. Available at <https://www.hsib.org.uk/investigations-and-reports/missed-detection-of-lung-cancer-on-chest-x-rays-of-patients-being-seen-in-primary-care/> (Accessed 22 December 2021).

Heit, J. A., Spencer, F. A. and White, R. H. (2016) The epidemiology of venous thromboembolism. *Journal of Thrombosis and Thrombolysis*, 41 (1), 3-14.
Helander, M. (2006) *A Guide to Human Factors and Ergonomics*, 2nd edition. Boca Raton, FL: CRC Taylor & Francis.

Hoffman, R. R., Ward, P., Feltovich, P. J., DiBello, L., Fiore, S. M. and Andrews, D. H. (2013) *Accelerated Expertise*. New York: Psychology Press.

Holden, R. J., Carayon, P., Gurses, A. P., Hoonakker, P., Hundt, A. S., Ozok, A. A. and Rivera-Rodriguez, A. J. (2013) SEIPS 2.0: a human factors framework for studying and improving the work of healthcare professionals and patients. *Ergonomics*, 56 (11), 1669-1686.

Hollnagel, E. (2009) *The ETTO Principle: Efficiency-Thoroughness Trade-Off. Why Things That Go Right Sometimes Go Wrong*. Farnham: Ashgate.

Howard, L. S. G. E., Barden, S., Condliffe, R., Connolly, V., Davies, C. W. H., Donaldson, J., Everett, B., Free, C., Horner, D., Hunter, L., Kaler, J., Nelson-Piercy, C., O-Dowd, E., Patel, R., Preston, W., Sheares, K. and Campbell, T. (2018) British Thoracic Society Guideline for the initial outpatient management of pulmonary embolism (PE). *Thorax*, 73 (Suppl 2), ii1-ii29.

Jiménez, D., García-Sánchez, A., Rali, P., Muriel, A., Bikdeli, B., Ruiz-Artacho, P., Le Mao, R., Rodríguez, C., Hunt, B.J. and Monreal, M. (2021) Incidence of VTE and bleeding among hospitalized patients with coronavirus disease 2019: a systematic review and meta-analysis. *Chest*, 159 (3), 1182-1196.

Joint Royal Colleges of Physicians Training Board. (n.d.) Internal medicine: about the specialty [Online]. Available at <https://www.jrcptb.org.uk/internal-medicine> (Accessed 29 November 2021).

Kaczmarczyk, J., Davidson, R., Bryden, D., Haselden, S. and Vivekananda-Schmidt, P. (2016) Learning decision making through serious games. *The Clinical Teacher*, 13 (4), 277-282.

Kahneman, D. (2012) *Thinking, Fast and Slow*. London: Penguin.

Kauppi, J. M., Airaksinen, K. E. J., Saha, J., Bondfolk, A., Pouru, J.-P., Purola, P., Jaakkola, S., Lehtonen, J., Vasankari, T., Juonala, M. and Kiviniemi, T. (2021) Adherence to risk-assessment protocols to guide computed tomography pulmonary angiography in patients with suspected pulmonary embolism. *European Heart Journal Quality of Care & Clinical Outcomes*, 16, qcab020.

Klein, G. (1989) Do decision biases explain too much? *Human Factors Society Bulletin*, 32 (5), 1-3.

Klein, G. (1997) The Recognition-Primed Decision (RPD) model: looking back, looking forward, in *Naturalistic Decision Making*, 1st edition. Mahwah, NJ: Lawrence Erlbaum Associates, Inc., 49-60.

Klein, G. (1999) *Sources of Power*. Cambridge, Mass., London: MIT Press.

Klein, G. (2008) Naturalistic decision making. *Human Factors*, 50 (3), 456-460.

Kline, J. A., Mitchell, A. M., Kabrhel, C., Richman, P. B. and Courtney, D. M. (2004) Clinical criteria to prevent unnecessary diagnostic testing in emergency department patients with suspected pulmonary embolism. *Journal of Thrombosis and Haemostasis*, 2 (8), 1247-1255.

Malas, M. B., Naazie, I. N., Elsayed, N., Mathlouthi, A., Marmor, R. and Clary, B. (2020) Thromboembolism risk of COVID-19 is high and associated with a higher risk of mortality: a systematic review and meta-analysis. *Eclinical Medicine*, 29, 100639.

Militello, L. and Hutton, R. J. (1998) Applied cognitive task analysis (ACTA): a practitioner's toolkit for understanding cognitive task demands. *Ergonomics*, 41 (11), 1618-1641.



Cognitive Task Analysis (ACTA) Methodology. Navy Personnel Research and Development Center, NPRDC-TN-98-4.

Moulton, C. and Mann, C. (2021) Emergency medicine: Getting It Right First Time [Online]. Available at <https://www.gettingitrightfirsttime.co.uk/medical-specialties/emergency-medicine/> (Accessed 14 December 2021).

Murray, D. J., Boyle, W. A., Beyatte, M. B., Knittel, J. G., Kerby, P. W., Woodhouse, J. and Boulet, J. R. (2018) Decision-making skills improve with critical care training: Using simulation to measure progress. *Journal of Critical Care*, 47, 133-138.

National Confidential Enquiry into Patient Outcome and Death. (2019) Know the score. A review of the quality of care provided to patients aged over 16 years with a new diagnosis of pulmonary embolism [Online]. Available at <https://www.ncepod.org.uk/2019pe.html> (Accessed 29 October 2020).

National Fire Chiefs Council. (n.d.a) Control measure – decision-making [Online]. Available at <https://www.ukfrs.com/guidance/search/decision-making> (Accessed 1 July 2021).

National Fire Chiefs Council. (n.d.b) Complementing incident command experience: a guide for fire and rescue [Online]. Available at <https://www.ukfrs.com/foundation-knowledge/complementing-incident-command-experience-guide-fire-and-rescue-services> (Accessed 8 July 2021).

National Institute for Health and Care Excellence. (2020b) Venous thromboembolic diseases: diagnosis, management and thrombophilia testing. NICE guideline [NG158] [Online]. Available at <https://www.nice.org.uk/guidance/ng158> (Accessed 24 February 2021).

National Institute for Health and Care Excellence. (2020c) Venous thromboembolic diseases: diagnosis, management and thrombophilia testing: evidence reviews [Online]. Available at <https://www.nice.org.uk/guidance/ng158/evidence/evidence-reviews-8710588333?tab=evidence> (Accessed 24 February 2021).

National Patient Safety Agency. (2010) Reducing treatment dose errors with low molecular weight heparins. Rapid response report NPSA/201/RRR014 [Online]. Available at <https://webarchive.nationalarchives.gov.uk/20171030124651/http://www.nrls.npsa.nhs.uk/resources/type/alerts/?entryid45=75208> (Accessed 24 February 2021).

NHS Digital. (n.d.) Hospital accident & emergency activity. Official statistics [Online]. Available at <https://digital.nhs.uk/data-and-information/publications/statistical/hospital-accident--emergency-activity> (Accessed 25 August 2021).

NHS England and NHS Improvement. (n.d.) A&E attendances and emergency admissions 2021-22 [Online]. Available at <https://www.england.nhs.uk/statistics/statistical-work-areas/ae-waiting-times-and-activity/ae-attendances-and-emergency-admissions-2021-22/> (Accessed 16 August 2021).

NHS England and NHS Improvement. (2019) The NHS Long Term Plan [Online]. Available at <https://www.longtermplan.nhs.uk/publication/nhs-long-term-plan/> (Accessed 16 August 2021).

NHS England and NHS Improvement. (2020) Transformation of urgent and emergency care: models of care and measurement [Online]. Available at <https://www.england.nhs.uk/wp-content/uploads/2020/12/transformation-of-urgent-and-emergency-care-models-of-care-and-measurement.pdf> (Accessed 16 August 2021).

NHS England and NHS Improvement. (2022) 2022/23 priorities and operational planning guidance [Online]. Available at <https://www.england.nhs.uk/wp-content/uploads/2021/12/B1160-2022-23-priorities-and-operational-planning-guidance-v2.pdf> (Accessed 11 February 2022).

O'Donovan, R. and McAuliffe, E. (2020) Exploring psychological safety in healthcare teams to inform the development of interventions: combining observational, survey and interview data. *BMC Health Services Research*, 20 (1), 810.

Patel, L., Gandhi, D., Westergard, E., Ornes, M., Lillyblad, M. and Skeik, N. (2021) COVID-19 and venous thromboembolism: known and unknown for imaging decisions. *World Journal of Radiology*, 13 (3), 64-74.

Patel, R., Sandars, J. and Carr, S. (2015) Clinical diagnostic decision-making in real life contexts: a trans-theoretical approach for teaching: AMEE Guide No. 95. *Medical Teacher*, 37 (3), 211-227.

Perera, M., Aggarwal, L., Scott, I. A. and Cocks, N. (2017) Underuse of risk assessment and overuse of computed tomography pulmonary angiography in patients with suspected pulmonary thromboembolism. *Internal Medicine Journal*, 47 (10), 1154-1160.



Phillips, J. K., Klein, G. and Sieck, W. R. (2004) Expertise in judgment and decision making: a case for training intuitive decision skills, in D. J. Koehler and N. Harvey (eds.), Blackwell Handbook of Judgment and Decision Making. Blackwell Publishing, 297-315.

Richards, M. (2020) Diagnostics: recovery and renewal. Report of the Independent Review of Diagnostic Services for NHS England [Online]. Available at <https://www.england.nhs.uk/wp-content/uploads/2020/11/diagnostics-recovery-and-renewal-independent-review-of-diagnostic-services-for-nhs-england-2.pdf> (Accessed 27 August 2021).

Richardson, S., Cohen, S., Khan, S., Zhang, M., Qiu, G., Oppenheim, M. I. and McGinn, T. (2020) Higher imaging yield when clinical decision support is used. Journal of the American College of Radiology, 17 (4), 496-503.

Rohacek, M., Buatsi, J., Szucs-Farkas, Z., Kleim, B., Zimmermann, H., Exadaktylos, A. and Stoupis, C. (2012) Requesting CT pulmonary angiography to exclude pulmonary embolism: defense versus evidence in the emergency room. Intensive Care Medicine, 38 (8), 1345-1351.

Royal College of Emergency Medicine. (2016) RCEM standard. Consultant sign-off [Online]. Available at https://res.cloudinary.com/studio-republic/images/v1635599020/Consultant_Sign_Off_Standard_June_2016/Consultant_Sign_Off_Standard_June_2016.pdf?_i=AA (Accessed 25 January 2022).

Royal College of Emergency Medicine. (2017) Delivering quality and models of care: initial assessment of ED patients [Online]. Available at <https://rcem.ac.uk/service-design-delivery-3/> (Accessed 24 February 2021).

Royal College of Emergency Medicine. (2019) Emergency care advanced clinical practitioner curriculum and assessment (adult only), version 2.0 [Online]. Available at https://rcem.ac.uk/wp-content/uploads/2021/10/EC_ACP_Curriculum_2017_Adult_Only-for_publication-Last_Edit_14-03-2019.pdf (Accessed 11 February 2022).

Royal College of Emergency Medicine. (2020) RCEM CARES during the coronavirus pandemic [Online]. Available at <https://rcem.ac.uk/rcem-cares/> (Accessed 4 August 2021).

Royal College of Emergency Medicine. (2021a) Generic Professional Capabilities Framework: RCEM curriculum [Online]. Available at <https://rcemcurriculum.co.uk/generic-professional-capabilities-framework/> (Accessed 10 June 2021).

Royal College of Emergency Medicine (2021b) RCEM acute insight series: crowding and its consequences [Online]. Available at https://rcem.ac.uk/wp-content/uploads/2021/11/Why_Emergency_Department_Crowding_Matters_v2.pdf#:~:text=Crowding%20is%20associated%20with%20increased%20mortality%20and%20increased,Against%20the%20backdrop%20of%20long%20ambulance%20delays%202 (Accessed 29 November 2021).

Royal College of Nursing. (n.d.) Revalidation requirements: reflection and reflective discussion [Online]. Available at <https://www.rcn.org.uk/professional-development/revalidation/reflection-and-reflective-discussion> (Accessed 24 June 2021).

Royal College of Physicians. (n.d.) Faculty of Physician Associates. Who are physician associates? [Online]. Available at <https://fparcp.co.uk/about-fpa/Who-are-physician-associates> (Accessed 11 February 2022).

Royal College of Physicians. (2021) Physician associate curriculum: draft [Online]. Available at https://www.gmc-uk.org/-/media/documents/draft---physician-associate-curriculum---15092021_pdf-87634359.pdf (Accessed 11 February 2022).

Royal College of Radiologists. (n.d.) About iRefer [Online]. Available at <https://www.rcr.ac.uk/clinical-radiology/being-consultant/rcr-referral-guidelines/about-irefer> (Accessed 28 January 2022).

Royal College of Radiologists. (2021) Appropriateness of usage of computed tomography pulmonary angiography (CTPA) investigation of suspected pulmonary embolism [Online]. Available at <https://www.rcr.ac.uk/audit/appropriateness-usage-computed-tomography-pulmonary-angiography-ctpa-investigation-suspected> (Accessed 11 June 2021).

Sharifzadeh, N., Kharrazi, H., Nazari, E., Tabesh, H., Edalati Khodabandeh, M., Heidari, S. and Tara, M. (2020) Health education serious games targeting health care providers, patients, and public health users: scoping review. *JMIR Serious Games*, 58 (1), e13459.

Stanton, N. (2013) *Human Factors Methods*. Farnham, UK: Ashgate Publishing Ltd.

Sullivan, E. J. and Garland, G. (2013) *Practical Leadership and Management in Healthcare: For Nurses and Allied Health Professionals*, 2nd edition. Harlow, England: Pearson.

Tholin, B., Ghanima, W., Einvik, G., Aarli, B., Brønstad, E., Skjønsberg, O.H. and Stavem, K. (2021) Incidence of thrombotic complications in hospitalised and non-hospitalised patients after COVID-19 diagnosis. *British Journal of Haematology*, 194 (3), 542-546.

Tran, A., Redley, M. and de Wit, K. (2021) The psychological impact of pulmonary embolism: a mixed-methods study. *Research and Practice in Thrombosis and Haemostasis*, 5 (2), 301-307.

van Belle, A., Büller, H. R., Huisman, M. V., Huisman, P. M., Kaasjager, K., Kamphuisen, P. W., Kramer, M. H. H., Kruip, M. J. H. A., Kwakkel-van Erp, J. M., Leebeek, F. W. G., Nijkeuter, M., Prins, M. H., Sohne, M. and Tick, L. W. (2006) Effectiveness of managing suspected pulmonary embolism using an algorithm combining clinical probability, D-dimer testing, and computed tomography. *JAMA*, 295 (2), 172-179.

Wang, R., DeMaria, S., Jr, Goldberg, A. and Katz, D. (2016). A systematic review of serious games in training health care professionals. *Simulation in Healthcare*, 11 (1), 41-51.

Wells, P. S., Anderson, D. R., Rodger, M., Stiell, I., Dreyer, J. F., Barnes, D., Forgie, M., Kovacs, G., Ward, J. and Kovacs, M. J. (2001) Excluding pulmonary embolism at the bedside without diagnostic imaging: management of patients with suspected pulmonary embolism presenting to the emergency department by using a simple clinical model and D-dimer. *Annals of Internal Medicine*, 135 (2), 98-107.

7 Appendices

7.1 Data extraction from the Strategic Executive Information System (StEIS)

Source:	StEIS
Date of extraction:	13 November 2020
Reported incident dates:	01 January 2020 - 31 October 2020
Filters:	N/A
Free text terms:	'pulmonary' 'embolism' 'emboli' 'PE'
Notes:	203 incidents identified and description field of each reviewed


7.2 Systems Engineering Initiative for Patient Safety (SEIPS)

SEIPS is a systems engineering approach with human factors principles embedded within it (**see figure 11**). SEIPS describes how components of the work system produce work processes which result in different outcomes. Work system factors are described below (Holden et al, 2013; Carayon et al, 2006):

- person(s): the people working in the particular system and the patient
- tasks: undertaken by the persons which may vary in complexity or variety
- tools and technology: used to undertake the tasks which may vary in usability and functionality
- internal environment: the physical space around the persons, for example layout, noise and temperature
- organisation: conditions external to the persons to support the organisation of, for example, resources and activity
- external environment: factors outside of the healthcare institution that might include policy, societal or economic factors.

Processes can be physical, cognitive, or behavioural and lead to outcomes for the patients, professionals or healthcare institutions. The interactions between the various components of the work system lead to different outcomes, both positive and negative. The framework includes feedback loops which represent the adjustments systems make over time.

Figure 11 Systems Engineering Initiative for Patient Safety (SEIPS) (Holden et al, 2013; Carayon et al, 2006)



7.3 Applied Cognitive Task Analysis – method

Overview of the method

An Applied Cognitive Task Analysis (ACTA) involves the following steps (Stanton, 2013):

- Observation of the task of interest with development of a task diagram with expert input.
- Identification of the steps in the task that require cognitive skills.
- Exploring of the cognitive aspects of the task where expertise is used, what those areas of expertise are and how they are used (termed the knowledge audit). Various probes are used to support this (see table 5).
- Identification of difficult cognitive elements, common errors and potential cues or strategies to overcome the difficulty.
- Simulation may also be used where experts are presented with typical scenarios and the steps they describe to manage it are examined from a cognitive perspective.

Table 5 Knowledge audit probes, adapted from Militello and Hutton (1998)

Past and future	Is there a time when you walked into a situation and knew exactly how things got there and where they were headed?
Big picture	What is important about the big picture for this task?
Noticing	Have you had experiences where a situation just ‘popped’ out at you?
Job smarts	When you do this task are there ways of working smartly that you have found useful?
Opportunities/ improvising	When have you improvised in this task or noticed opportunities to do things better?
Self monitoring	Is there a time when you needed to change the way you were performing in order to get the job done?
Anomalies	Have there been times when you spotted a deviation or knew something was not right?
Equipment difficulties	Have there been times that equipment pointed you in the wrong direction?
Scenario from hell	If you were going to devise a scenario to really show someone what this task was all about, what would you put in it?



The investigation's ACTA

The investigation's ACTA approach was:

- Training of the investigation team by a subject matter advisor.
- Identification of cognitive elements following observations in two emergency departments (EDs).
- Development of a cognitive task diagram.
- Validation of the cognitive task diagram at the previously visited EDs.
- Knowledge audits with experts and novices across six EDs.
- Discussion of scenarios and their challenging cognitive elements.
- Development of a cognitive demands table and presentation of findings.

7.4 Applied Cognitive Task Analysis - findings

The following sections provide supporting information to **section 4.2** in the main report.

Initial assessment

What do experts do?

The investigation was told that expert nursing staff:

- recognise that early focus on a diagnosis, such as PE, may influence later decisions.
- seek to identify symptoms to guide the selection of tests that may later support diagnosis.
- recognise that a D-dimer test has a specific use, and only request one following expert consultation.

[Nurse] described never ordering a D-dimer at triage unless asked to by a consultant. They recalled diagnoses being missed because PE had been fixed on early. They sent a coagulation [blood] sample to which a D-dimer could be later added.

Band 7 Nurse



Why is it difficult for novices?

The investigation was told by novice staff that initial assessment is difficult for novice nurses because:

- They are not trained to assess symptoms and select tests; instead training focuses on triage.
- It is seen as an isolated process without recognition of its influence on later diagnosis and treatment.
- They rarely receive feedback on whether a D-dimer test was appropriate, and do not know the outcome of their patients.
- Blood order sets (requesting forms with pre-determined blood tests depending on the patient's symptoms) and strict procedures lead to automatic behaviours without consideration of implications.
- Because of the national operating standard, ED processes are performance focused rather than outcome focused.

[Nurse] at triage described that they ordered a D-dimer for any patient short of breath and with chest pain. This was to prevent 'breaching' of the 4-hour target having been previously told of the need to meet the target.

Band 6 Nurse

Further clinical assessment

What do experts do?

The investigation was told that expert staff (consultants/senior doctors/experienced advanced clinical practitioners):

- Recognise that they can never be 100% sure of a diagnosis without relevant tests, but will have a comfort threshold for when they feel testing is needed based on experience.
- Use discriminatory questioning to account for each symptom, sign and investigation result. Any left unexplained require further investigation. For example, unexplained tachycardia (fast heart rate).
- Recognise that observations and investigation results must be interpreted with knowledge of the individual patient and what is normal for them.



[Doctor] reviewed a patient prior to discharge. The patient had presented with chest pain and had a normal troponin. Their heart rate was still elevated compared to their normal. A subsequent D-dimer was positive and a computerised tomography pulmonary angiography (CTPA) showed PE.

Consultant

- Recognise the potential for an undiagnosed and significant condition in patients with non-specific symptoms and risk factors. They use 'gut feeling'.
- Look for subtle clues to support their understanding of whether a condition is present.
- Recognise that well patients can still have a significant condition.

[Practitioner] was observed to get a mildly breathless patient up and walking; this identified a drop in their oxygen levels on exertion not identified at rest. Further tests suggested a PE.

Advanced Clinical Practitioner

- Appreciate current context, for example the COVID-19 pandemic, on patients' symptoms.
- Understand, recognise and adapt to the constraints of the healthcare system, such as CTPA access and the national operating standard.
- Appreciate the specific role of D-dimers in the diagnostic pathway for PE and are comfortable following it.

[Doctor] described requesting a D-dimer only when they had first reviewed the patient and considered a PE. They recalled a patient with temperature, shortness of breath, raised D-dimer and evidence of pneumonia on the X-ray. They accounted for the D-dimer rise by the infection.

Consultant

- Seek feedback on outcomes of the patients they have treated to reflect on their decisions, and learn from when they were wrong.
- Discuss with fellow experts when they are uncertain about what decision to make to access multiple opinions.

Why is it difficult for novices?

The investigation was told that further assessment is difficult for novice staff because:

- They may see few or no PEs or aortic dissections in their career.
- Training focuses on textbook symptoms and signs suggestive of PE and aortic dissection, leading to rule-based decisions.
- They may reassure themselves that a significant illness/condition is not present in patients who do not fit the rules or who seem well.

[Doctor] saw a patient with high blood pressure and chest pain through to their back. As the patient looked well they discounted an aortic dissection. A consultant review prior to discharge resulted in further tests and diagnosis of an aortic dissection.

Non-consultant doctor

- Training focuses on common conditions and so novices may not expect young, fit and well people to have a PE.

[Student] described that they had been taught that young adults with tachycardia and anxiety are probably suffering from panic attacks. They therefore did not expect younger adults to have a PE.

Medical student

- Limited experience results in fitting symptoms, signs and investigation results to a 'best fit' diagnosis without accounting for all unusual test results, symptoms or signs.

[Doctor] saw a patient with shortness of breath. They noted the patient had asthma and decided this the likely diagnosis despite a clear chest and unexplained cause for the shortness of breath.

Non-consultant doctor

- Limited experience may also mean higher confidence in a diagnosis having not seen the variable signs and symptoms that someone with a condition may have.



- Documentation of suspected diagnoses on referral letters and initial assessment documents, and what tests have already been ordered, may cue a novice to consider a specific diagnosis to the detriment of another.
- Less experienced staff may require reassurance for their decisions which may mean ordering tests such as D-dimer outside of their intended use.
- Clinical demand limits opportunities for structured case reviews with experts. Experts may only have capacity to discuss patients, rather than see them in person with the novice.
- Staff may be worried about what others think of them and negative responses to previous decisions may alter future decisions.

[Doctor] described pushback from radiology when requesting CTPAs from ED. They therefore would only request one if they had a high suspicion of PE.

Non-consultant doctor

General practitioner perspective

ED staff told the investigation that their assessments were often supported by the availability of tests and several hours to observe patients. In contrast they described that GPs in primary care have short appointments within which to make decisions and limited access to tests. To examine the primary care perspective, the investigation approached an experienced GP.

The GP described various cues, beyond standard history and examination, that developed his suspicion of a significant medical problem. He walks to the waiting room to collect patients providing early information through cues such as social interaction and the effects of exertion when walking. He described how he “teases out” the potential causes behind symptoms and signs, and is suspicious of those without explanation. He spoke of being concerned by the patient who “is not quite right”. Examples included a young, fit adult with lower than usual oxygen saturations, or an active person who is still able to exercise, but not as intensely.

The GP shared that with experience comes learning from unintended outcomes; these shape levels of perceived risk and make clinicians suspect conditions such as PE. As an experienced practitioner, he recognised the importance of “gut instinct” and that as he has become more experienced, he has felt less inhibited to trust it. This has also meant he has become less worried about what others think as long as he is doing what he feels is best for the patient.



A novice may be worried about what others think and may fit unexplained symptoms and signs to a diagnosis. Novices may not tease out the specifics of a patient's signs and symptoms, focusing on common conditions and accepting them at face value.

Finally, the GP described the context within which his decisions are made. Telephone consultations have become more common, but have removed visual cues. There is also significant demand for GP services, making decisions within short appointments difficult.



WWW.HSIB.ORG.UK

 [@hsib_org](https://twitter.com/hsib_org)



HEALTHCARE SAFETY
INVESTIGATION BRANCH

Further information

More information about HSIB – including its team, investigations and history – is available at www.hsib.org.uk

If you would like to request an investigation then please read our **guidance** before contacting us.

 [@hsib_org](https://twitter.com/hsib_org) is our Twitter handle. We use this feed to raise awareness of our work and to direct followers to our publications, news and events.

Contact us

If you would like a response to a query or concern please contact us via email using enquiries@hsib.org.uk

We monitor this inbox during normal office hours - Monday to Friday from 09:00 hours to 17:00 hours. We aim to respond to enquiries within five working days.

To access this document in a different format – including braille, large-print or easy-read – please contact enquiries@hsib.org.uk

© Healthcare Safety Investigation Branch copyright 2022. Any enquiries regarding this publication should be sent to us at enquiries@hsib.org.uk